


Name: KEY

Chemistry 20

Worksheets



Worksheet 1.1: Atomic Structure

1. Complete the following table. Using symbols, provide an example of each category using the element provided. For the average atom assume that the mass number is the atomic mass rounded off. The first one is done. (8 marks) (1/2 mark off for each mistake in each row).

	Average Atom	Monoatomic Polyatomic or diatomic element	Most common simple Ion	Complex or polyatomic ion	Isotope of the average	Ionic compound	Molecular compound	Acid compound
chromium	^{52}Cr	Cr	Cr^{3+}	CrO_4^{2-}	^{53}Cr	Cr_2O_3	NONE	NONE
sulphur	^{32}S	$\text{S}_{8(s)}$	S^{2-}	SO_4^{2-} or SO_3^{2-}	^{33}S	$\text{Na}_2\text{S}_{(aq)}$	$\text{SO}_{2(g)}$	$\text{H}_2\text{SO}_{4(aq)}$ $\text{H}_2\text{SO}_{3(aq)}$
nitrogen	^{14}N	$\text{N}_{2(g)}$	N^{3-}	NO_3^- or NO_2^-	^{13}N	$\text{Na}_3\text{N}_{(aq)}$	$\text{NO}_{2(g)}$	$\text{HNO}_{3(aq)}$ $\text{HNO}_{2(aq)}$

2. Complete the following table. Note that the mass number can change for isotopes. If there is not a noble gas with the same # of electrons, than put NONE. The first one is done.(6 marks) (1/2 mark off for each mistake in each row)

Atom or Ion name	Atom or ion symbol	Atomic number	Mass number	Protons	Electrons	Neutrons	Noble Gas with same # of electrons
sodium ion	$^{23}\text{Na}^+$	11	23	11	10	12	neon
aluminum atom	^{28}Al	13	28	13	13	15	neon
chloride ion	Cl^{1-}	17	34	17	18	17	argon

Worksheet 1.2: Compounds

Complete the following table (Assume water is used with ionic compounds): (16 marks)


I = M (NH ₄) + N M = non metals A = H (acid) pg 8/9 in databook	chemical formula(add states)	chemical name
Ionic	K ₂ SO _{3(aq)}	Metal name + polyatomic name potassium sulfite
ACID	H ₂ SO _{4(aq)}	sulphuric acid
IONIC	Na ₂ S ₂ O ₃ •1 H ₂ O _(s)	sodium thiosulfate monohydrate
MOLECULAR	C ₂ H ₅ OH(l)	ethanol
IONIC	Pb ^{2+/4+} (SO ₄) ₂ ²⁻ _(aq) 4+ = 4-	lead (IV) sulphate
MOLECULAR	P ₅ O ₁₀ (g)	Pentaphosphorus dexaoxide
MOLECULAR	C₁₂H₂₂O₁₁ (s)	sucrose
IONIC	Na₂SiO₃(aq)	sodium silicate
MOLECULAR	NH ₃ (g)	Ammonia is not ammonium

MOLECULAR (NOT AN ACID)	$\text{H}_2\text{O}_2(l)$	Hydrogen peroxide
MOLECULAR	$\text{SO}_3(g)$ does not equal $\text{SO}_3^{2-}(aq)$	Sulfur trioxide (NOT sulfite)
IONIC (NH_4^+ is an ion)	$(\text{NH}_4)_3\text{PO}_4(aq)$	ammonium phosphate
IONIC	Cu^{2+} SO_4^{2-} $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}(s)$	copper (II) sulphate pentahydrate
MOLECULAR (MEMORIZED)	$\text{C}_3\text{H}_8(g)$	propane
ACID	$\text{CH}_3\text{COOH}(aq)$	ethanoic acid
MOLECULAR	$\text{O}_3(g)$	OZONE
MOLECULAR	$\text{HOH}_{(g,l,s)}$	water

Worksheet 1.3: Reactions

Complete the following reactions, identify the reaction type and balance the equation. (3 marks each; 15 marks total)

- 1) mercury (II) oxide is broken down into its elements by heating.



- 2) a nickel strip is placed in a gold (III) sulfate solution



- 3) phosphoric acid reacts with iron (III) oxide.



$$\text{H} = 3 \times 6$$

$$\text{H} = 2 \times 2$$

$$\text{O} = 3$$

$$\text{O} = 4 \times 3$$


$$\text{PO}_4 = 1 \times 2$$

$$\text{PO}_4 = 4 \times 2$$


$$\text{Fe} = 2$$

$$\text{Fe} = 1 \times 2$$

- 4) butane is burned in air



$$2? = 8 + 5; ? = 6.5$$



- 5) sulfur combines with oxygen to form sulfur trioxide



Worksheet 1.4: Mole Problems

1. What is the molar mass of hydrogen peroxide?




$$\text{H} \times 2 = 2.02$$

$$\text{O} \times 2 = 32.00$$

$$\text{Total} = 34.02 \text{ g/mol}$$

2. What is the molar mass of lead (II) nitrate?



$$\text{Pb} \times 1 = 207.2$$

$$\text{N} \times 2 = 28.02$$

$$\text{O} \times 6 = 96.00$$

$$\text{TOTAL} = 231.22 \text{ g/mol}$$

3. How many moles in 30.6 g of copper?

$$\text{Step 1) } n=?; m = 30.6\text{g}; M = 63.55\text{g/mol}$$

$$\text{Step 2) } n = m/M$$

$$\text{Step 3) } n = 30.6\text{g}/63.55\text{g/mol}$$

$$\text{Step 4) } n = 0.481510\dots \text{ mol}; 0.482 \text{ mol or } 482 \text{ mmol} \text{ (3 significant digits; divide by E-3 to get mmol)}$$

4. How many moles in 6.55×10^{19} atoms of zinc?

$$\text{Step 1) } n=?; p = 6.55\text{E}19 \text{ atoms}; P = 6.02\text{E}23 \text{ atoms/mol}$$

$$\text{Step 2) } n = p/P$$

$$\text{Step 3) } n = 6.55\text{E}19 \text{ atoms} / 6.02\text{E}23 \text{ atoms/mol}$$

$$\text{Step 4) } n = 0.0001088039\dots \text{ mol}; 0.000109 \text{ mol or } 1.09 \times 10^{-4} \text{ mol or } 0.109 \text{ mmol or } 109 \text{ umol}$$

5. Determine the number of moles in 33.6 L of methane at STP?

$$\text{Step 1) } n=?; v = 33.6\text{L}; V = 22.4\text{L/mol}$$

$$\text{Step 2) } n = v/V$$

$$\text{Step 3) } n = 33.6\text{g}/22.4\text{L/mol}$$

$$\text{Step 4) } n = 1.50 \text{ mol} \text{ (3 significant digits)}$$

6. How many formula units in 3.99 mol of potassium carbonate? (K_2CO_3)
Step 1) $p=?$; $n = 3.99\text{mol}$; $P = 6.02 \text{E}23$ formula units/mol
Step 2) $p = n \times P$
- Step 3) $p = 3.99\text{mol} \times 6.02\text{E}23$ formula units/mol
- Step 4) $p = 2.40198 \text{E}24$ formula units; **$2.40\text{E}24$ formula units or 2.40×10^{24} formula units**
7. What volume of gas would be present in 0.955 mol at STP?; SATP?
Step 1) $v=?$; $n = 0.955$ mol; $V = 22.4\text{L/mol}$
- Step 2) $v = nV$
- Step 3) $v = 0.955\text{mol} \times 22.4 \text{L/mol}$ (STP) $v = 0.955\text{mol} \times 24.8\text{L/mol}$
- Step 4) $v = 21.392 \text{L}$; **21.4L** of gas at STP $v = 23.684$; **23.7L** of gas at SATP
8. What is the mass of 2.3 mol of carbon dioxide at STP?
Step 1) $n=?$; $m = 30.6\text{g}$; $M =$
C = $12.01 \times 1 = 12.01$
O = $16.00 \times 2 = 32.00$
TOTAL = 48.01g/mol
- Step 2) $m = nM$
- Step 3) $m = 2.3\text{mol} \times 48.01\text{g/mol}$
- Step 4) $m = 110.423 \text{g}$; **$1.1 \times 10^2 \text{g}$ or 0.11kg** (divide by $\text{E}3$ to change to kmol)
- NOTE: You have to change to scientific notation since you need 2 sig digs and 100 has 3/
9. How many moles in 100 mL of carbon dioxide at SATP?
Step 1) $n=?$; $v = 100\text{mL}$ OR 0.100L ; $V = 24.8\text{L/mol}$
- Step 2) $n = v/V$
- Step 3) $n = 0.100 \text{L} / 24.8 \text{L/mol}$
- Step 4) $n = 0.004032258\dots \text{mol}$; **0.00403mol or $4.03 \times 10^{-3} \text{mol}$ or 4.03mmol**
10. What volume of nitrogen monoxide would be present in 2.7 mol if the temperature is 25°C and the pressure is 100kPa ?
Step 1) $v=?$; $n = 2.7 \text{mol}$; $V = 24.8\text{L/mol}$ (SATP conditions)
- Step 2) $v = nV$
- Step 3) $v = 2.7 \text{mol} \times 24.8\text{L/mol}$
- Step 4) $v = 66.96 \text{L}$; **67L** (2 significant digits)

Worksheet 1.5: More difficult mole problems


- How many molecules in 2.00 mol of sulphur dioxide?
 $p=nP$; $p=2.00 \text{ mol} \times 6.02\text{E}23 \text{ molecules/mol}$
 $p= 1.20\text{E}24 \text{ molecules of sulphur dioxide}$
- How many molecules in 7.5 mol of chlorine?
 $p=nP$; $p=7.5 \text{ mol} \times 6.02\text{E}23 \text{ molecules/mol}$
 $p= 4.5\text{E}25 \text{ molecules of chlorine}$
- How many atoms of **copper** are in 0.088 mol of copper (I) oxide?
 $p=nP$; $p=0.088 \text{ mol} \times 6.02\text{E}23 \text{ molecules/mol}$
 $p= 5.2976\text{E}22 \text{ molecules of Cu}_2\text{O}$
 Cu_2O ; $p_{\text{Cu}}= 2 \times = 1.059\text{E}23$; **$1.1\text{E}23 \text{ atoms of copper}$**
- How many mol of **magnesium ions** are in 1.00×10^{20} formula units of magnesium nitride?
 $n=p$; $n=1.00\text{E}20 \text{ formula units}$; $n= 1.66 \times 10^{-4} \text{ mol or } 0.166 \text{ mmol}$
P $6.02\text{E}23 \text{ formula units/mol}$
NOT DONE YET....
 Mg_3N_2 ; $n_{\text{Mg}}= 3 \times 0.166 \text{ mmol} = 0.498 \text{ mmol}$
- What is the mass of 14.6 L of carbon monoxide at STP?
C = 12.01 $n=v/V$; $n=14.6\text{L}/22.4\text{L/mol}$; $n=0.651785714\dots \text{ mol}(\text{don't round})$
O = 16.00 $m=nM$; $m=0.651785714\dots\text{mol} \times 28.01\text{g/mol}$; $m= 18.256\dots \text{ g}$; **18.3g**
28.01g/mol

6. How many atoms of xenon are in 15 L at SATP?
 $n = v/V$; $n = 15\text{L}/24.8\text{L/mol}$; $n = 0.6048387\dots\text{mol}$
 $p = nP$; $p = 0.6048387\dots\text{mol} \times 6.02\text{E}23 \text{ atoms/mol}$; $p = 3.6411\dots\text{E}23$
 $p = 3.6 \text{ E } 23 \text{ atoms}$
7. How many moles of **carbon and oxygen** are in 6.02×10^{23} molecules of carbon dioxide?
 $n = p$; $n = \frac{6.02\text{E}23 \text{ formula units}}{6.02\text{E}23 \text{ formula units/mol}}$; $n = 1.00 \text{ mol}$
NOT DONE YET....
 CO_2 ; $n_{\text{C}} = 1 \times 1.00 = 1.00 \text{ mol of carbon}$; $n_{\text{O}} = 2 \times 1.00 = 2.00 \text{ mol of oxygen}$
8. When studying reactions what unit are most quantities converted into? (HINT: What do the coefficients in front of a balanced equation represent?)
Most quantities are converted into moles. The coefficients represent moles
9. What are the temperature, pressure and molar volume of a gas at STP?
 $T = 0\text{C} (273.15\text{K})$; $P = 101.325 \text{ kPa}$; $V = 22.4 \text{ L/mol}$
10. What are the temperature, pressure and molar volume of a gas at SATP?
 $T = 25 \text{ C} (298.15 \text{ K})$; $P = 100 \text{ kPa}$; $V = 24.8 \text{ L/mol}$

Worksheet 2.1: Mole to Mole Stoichiometry


Directions: Write balanced equations with states. Solve the problem. Assume water is available.

1. Liquid water decomposes into its elements. How many moles of hydrogen gas are produced if 0.500 mol of water is used?




$$\text{step 3) } \frac{0.500 \text{ mol of H}_2\text{O}_{(g)} \times 2 \text{ mol of H}_{2(g)}}{2 \text{ mol of H}_2\text{O}_{(g)}} = \mathbf{0.500 \text{ mol of H}_2(g)}$$

2. Sulphur reacts with barium oxide. How many moles of sulphur are needed if 2.00 mol of barium oxide is used?




$$\text{3) } \frac{2.00 \text{ mol of BaO}_{(s)} \times 1 \text{ mol of S}_{8(s)}}{8 \text{ mol of BaO}_{(s)}} = \mathbf{0.250 \text{ mol of S}_{8(s)}}$$

3. Methane gas burns. How many moles of oxygen gas are needed to completely burn 3.00 mol of methane?




$$\text{3) } \frac{3.00 \text{ mol of CH}_{4(g)} \times 2 \text{ mol of O}_{2(g)}}{1 \text{ mol of CH}_{4(g)}} = \mathbf{6.00 \text{ moles of O}_{2(g)}}$$

4. Sodium and phosphorus react. How many moles of phosphorus are needed if 0.600 mol of sodium metal is used?




$$\text{3) } \frac{0.600 \text{ mol of Na}_{(s)} \times 1 \text{ mole of P}_{4(s)}}{12 \text{ moles of Na}_{(s)}} = \mathbf{0.0500 \text{ moles of P}_{4(s)}}$$

5. Magnesium phosphate reacts with lithium carbonate. How many moles of lithium carbonate are needed when 1.50 mol of magnesium phosphate is used?




$$\text{3) } \frac{1.50 \text{ mol of Mg}_3(\text{PO}_4)_{2(s)} \times 3 \text{ mol of Li}_2\text{CO}_{3(aq)}}{1 \text{ mol of Mg}_3(\text{PO}_4)_{2(s)}} = \mathbf{4.50 \text{ mol of Li}_2\text{CO}_{3(aq)}}$$

6. Sulphur dioxide decomposes. How many moles of sulphur dioxide are needed to produce 0.30 mol of sulphur?




$$3) \frac{0.30 \text{ mol of S}_{8(s)} \times 8 \text{ mol of SO}_{2(g)}}{1 \text{ mol of S}_{8(s)}} = 2.4 \text{ mol of Sulphur dioxide}$$

7. Magnesium chloride reacts with sodium. How many moles of sodium are needed to react with 0.0250 mol of magnesium chloride?




$$3) \frac{0.0250 \text{ mol MgCl}_{2(aq)} \times 2 \text{ mol of Na}_{(s)}}{1 \text{ mol of MgCl}_{2(aq)}} = 0.0500 \text{ mol of Na}_{(s)} (5.00 \times 10^{-2} \text{ mol})$$

8. Iron (II) phosphate reacts with tin (IV) nitride. How many moles of tin (IV) nitride are needed to produce 0.500 mol of iron (II) nitride?




$$3) \frac{0.500 \text{ mol of Fe}_3\text{N}_{2(s)} \times 1 \text{ mol of Sn}_3\text{N}_{2(s)}}{2 \text{ mol of Fe}_3\text{N}_{2(s)}} = 0.250 \text{ mol of tin (IV) phosphate}$$

9. Gasoline ($\text{C}_8\text{H}_{18(l)}$) is burned. How many moles of carbon dioxide are produced when 3.00 mol of gasoline is reacted?



$$3) \frac{3.00 \text{ mol of C}_8\text{H}_{18(l)} \times 8 (16) \text{ mol of CO}_{2(g)}}{1 (2) \text{ mol of C}_8\text{H}_{18(l)}} = 24.0 \text{ mol of carbon dioxide.}$$

10. Chlorine reacts with potassium bromide. How many moles of chlorine would be needed to completely use up 25 mol of potassium bromide?




$$3) \frac{25 \text{ mol of KBr}_{(aq)} \times 1 \text{ mol of Cl}_{2(g)}}{2 \text{ mol of KBr}_{(aq)}} = 13 \text{ mol of chlorine}$$

Worksheet 2.2: Mole to Quantity Stoichiometry

Directions: Solve the following hypothetical stoichiometry problems. Assume water is available.

1. When 6.5 mol of solid potassium chlorate breaks into solid potassium chloride and oxygen gas, what mass of solid potassium chloride is produced?



LINEAR METHOD

$$n = \frac{6.5 \text{ mol} \times 2 \text{ mol of KCl}_{(s)}}{2 \text{ mol of KClO}_{3(s)}} \times \frac{74.55 \text{ g of KCl}_{(s)}}{1 \text{ mol of KCl}_{(s)}} = 484 \text{ g} = 4.8 \times 10^2 \text{ g of KCl}_{(s)}$$

STEP BY STEP METHOD


2) no conversion

3) mol ratio: $n_R = n_G \times R/G$

$$6.5 \text{ mol of KClO}_{3(s)} \times 2 \text{ mol of KCl}_{(s)} / 2 \text{ mol of KClO}_{3(s)} = 6.5 \text{ moles of KCl}_{(s)}$$

4) $m = nM$ $m = 74.55 \text{ g/mol} \times 6.5 \text{ mol} = 484 \text{ g} = 4.8 \times 10^2 \text{ g of KCl}_{(s)}$

2. When 5.00 mol of methane burns, what volume of carbon dioxide at STP, will be produced?



LINEAR METHOD:


$$\frac{5.00 \text{ mol of CH}_4}{1 \text{ mol of CH}_4} \times \frac{1 \text{ mol of CO}_2}{1 \text{ mol of CH}_4} = \frac{22.4 \text{ L of CO}_2}{1 \text{ mol of CO}_2} = 112 \text{ L of CO}_2$$

STEP BY STEP METHOD: 2) no conversion

3) $n_R = n_G \times R/G = 5.00 \text{ mol of CH}_4 \times 1 \text{ mol of CO}_2 / 1 \text{ mol of CH}_4 = 5.00 \text{ mol of CO}_2$

4) $v = nV$ $v = 5.00 \text{ mol of CO}_2 \times 22.4 \text{ L/mol} = 112 \text{ L of CO}_2$

3. How many particles of hydrochloric acid is needed to neutralize 2.50 mol of calcium hydroxide?



LINEAR METHOD:

$$n = \frac{2.5 \text{ mol of Ca(OH)}_{2(s)} \times 2 \text{ mol of HCl}_{(aq)}}{1 \text{ mol of Ca(OH)}_{2(s)}} \times \frac{6.02 \times 10^{23} \text{ particles of HCl}_{(aq)}}{1 \text{ mol of HCl}_{(aq)}} = 3.01 \text{ E } 24 \text{ particles}$$

STEP BY STEP:


2) no conversion

3) $n_R = n_G \times R/G = \frac{2.5 \text{ mol of Ca(OH)}_{2(s)} \times 2 \text{ mol of HCl}_{(aq)}}{1 \text{ mol of Ca(OH)}_{2(s)}}$

$n = 5.0 \text{ mol}$

4) $p = n \times P$ $p = 6.02 \times 10^{23} \times 5.0 \text{ mol} = 3.01 \text{ E } 24 \text{ or } 3.01 \times 10^{24} \text{ particles of HCl}_{(aq)}$

4. When 5.25 mol of butane ($\text{C}_4\text{H}_{10(l)}$) burns, what volume of water vapour will be produced at SATP?



LINEAR METHOD:


$$\frac{5.25 \text{ mol of C}_4\text{H}_{10(l)}}{1 \text{ mol of C}_4\text{H}_{10(l)}} \times \frac{5 \text{ mol of H}_2\text{O}_{(g)}}{1 \text{ mol of C}_4\text{H}_{10(l)}} = \frac{24.8 \text{ L of H}_2\text{O}_{(g)}}{1 \text{ mol of H}_2\text{O}_{(g)}} = 651 \text{ L of H}_2\text{O}$$

STEP BY STEP METHOD: 2) no conversion

3) $n_R = n_G \times R/G = 5.25 \text{ mol of C}_4\text{H}_{10(l)} \times 5 \text{ mol of H}_2\text{O}_{(g)} / 1 \text{ mol of C}_4\text{H}_{10(l)} = 16.25 \text{ mol of H}_2\text{O}_{(g)}$

4) $v = nV$ $v = 16.25 \text{ mol of H}_2\text{O} \times 24.8 \text{ L/mol} = 651 \text{ L of water}$

5. When excess silver reacts with 3.45 mol of zinc phosphate, what mass of silver phosphate would be produced?



LINEAR METHOD:


$$\frac{3.45 \text{ mol of Zn}_3(\text{PO}_4)_2 \times 2 \text{ mol of } 2\text{Ag}_3\text{PO}_4 \times 418.58 \text{ g of Ag}_3\text{PO}_4}{1 \text{ mol of Zn}_3(\text{PO}_4)_2 \quad 1 \text{ mol of Ag}_3\text{PO}_4} = 2888 \text{ g} = \mathbf{2.89 \text{ kg}}$$

STEP BY STEP METHOD: 2) no conversion

$$3) \quad n_R = n_G \times R/G = \frac{3.45 \text{ mol of Zn}_3(\text{PO}_4)_2 \times 2 \text{ mol of } 2\text{Ag}_3\text{PO}_4}{1 \text{ mol of Zn}_3(\text{PO}_4)_2} = 6.90 \text{ mol}$$

$$4) \quad m = Mn = (418.58 \text{ g/mol})(6.90 \text{ mol}) = 2888 \text{ g} = \mathbf{2.89 \times 10^3 \text{ g or } 2.89 \text{ kg}}$$

6. When 3.00 mol of iron (II) hydroxide reacts with cobalt (II) phosphate, what mass of cobalt (II) phosphate is needed? G R



LINEAR METHOD:

$$\frac{3.00 \text{ mol of Fe}(\text{OH})_{2(aq)} \times 3 \text{ mol of Co}(\text{OH})_{2(s)} \times 552.55 \text{ g of Co}_3(\text{PO}_4)_{2(s)}}{3 \text{ mol of Fe}(\text{OH})_{2(aq)} \quad 1 \text{ mol of Co}_3(\text{PO}_4)_{2(s)}} = \mathbf{278.85 = 279 \text{ g of Co}(\text{OH})_2}$$


STEP BY STEP METHOD:

2) no conversion

$$3) \quad n_R = n_G \times R/G = \frac{3.00 \text{ mol of Fe}(\text{OH})_{2(aq)} \times 3 \text{ mol of Co}(\text{OH})_{2(s)}}{3 \text{ mol of Fe}(\text{OH})_{2(aq)}} = 3.00 \text{ mol of Co}(\text{OH})_{2(s)}$$

$$4) \quad m = Mn = (92.95 \text{ g/mol})(3.00 \text{ mol}) = \mathbf{278.85 = 279 \text{ g of Co}(\text{OH})_2}$$

7. In a neutralization reaction, 4.56 mol of sodium hydroxide neutralizes the sulphuric acid. What mass of water is produced? G R



LINEAR METHOD:

$$\frac{4.56 \text{ mol of NaOH}_{(aq)} \times 2 \text{ mol of H}_2\text{O}_{(g)} \times 18.02 \text{ g of H}_2\text{O}_{(g)}}{2 \text{ mol of NaOH}_{(aq)} \quad 1 \text{ mol of H}_2\text{O}_{(g)}} = \mathbf{8.22 \times 10^1 \text{ g of water}}$$


STEP BY STEP METHOD:

2) no conversion

$$3) \quad n_R = n_G \times R/G = \frac{4.56 \text{ mol of NaOH} \times 2 \text{ mol of H}_2\text{O}}{2 \text{ mol of NaOH}} = 4.56 \text{ mol of H}_2\text{O}$$

$$4) \quad m = Mn = (18.02 \text{ g/mol})(4.56 \text{ mol}) = \mathbf{8.22 \times 10^1 \text{ g of water}}$$

8. Hydrogen and 2.5 mol of nitrogen react to form ammonia. How many moles of ammonia will be produced at STP? SATP? G R



LINEAR METHODE

$$\frac{2.5 \text{ mol of N}_{2(g)} \times 2 \text{ mol of NH}_{3(g)} \times 22.4 \text{ L (24.8 L) of NH}_{3(g)}}{1 \text{ mol of N}_{2(g)} \quad 1 \text{ mol of NH}_{3(g)}} = (112) \mathbf{1.1E2 \text{ L of N}_{2(g)}}; (124) \mathbf{1.2E2 \text{ L of N}_{2(g)}}$$

STEP BY STEP METHOD:

2) no conversion


$$3) \quad n_R = n_G \times R/G = \frac{2.5 \text{ mol of N}_2 \times 2 \text{ mol of H}_2\text{O}}{1 \text{ mol of NaOH}} = 5.00 \text{ mol of N}_2$$

$$4) \quad v = nV = (5.00 \text{ mol})(22.4 \text{ L/mol}) = \mathbf{1.1E2 \text{ L of N}_2 \text{ STP}}; v = nV = (5.00 \text{ mol})(24.8 \text{ L/mol}) = \mathbf{1.2E2 \text{ L of N}_2 \text{ SATP}}$$

Worksheet 2.3: Quantity to Mole Stoichiometry

Directions: Solve the following hypothetical stoichiometry problems. Assume water is available.

1. How many moles of iron (III) oxide is produced when 5.6 g of iron burns with oxygen gas?



LINEAR METHOD:


$$\frac{5.6 \text{ g of Fe}_{(s)} \times 1 \text{ mol of Fe}_{(s)} \times 1 \text{ mol of Fe}_2\text{O}_{3(s)}}{55.85 \text{ g of Fe}_{(s)} \quad 2 \text{ mol of Fe}_{(s)}} = 0.050 \text{ mol of Fe}_2\text{O}_{3(s)}$$

STEP BY STEP METHOD:

2) $n = m/M = 5.6 \text{ g} / 55.85 \text{ g/mol} = 0.10 \text{ mol}$

3) $n_R = n_G \times R/G = 0.10 \text{ mol of Fe}_{(s)} \times \frac{1 \text{ mol of Fe}_2\text{O}_{3(s)}}{2 \text{ mol of Fe}_{(s)}} = 0.050 \text{ mol of Fe}_2\text{O}_{3(s)}$

2. When 4.00×10^{23} particles of methanol is burned, how many moles of water vapour are produced?



LINEAR METHOD:


$$\frac{4.00 \times 10^{23} \text{ part of CH}_3\text{OH}_{(l)} \times 1 \text{ mol of CH}_3\text{OH}_{(l)} \times 2 \text{ mol of H}_2\text{O}_{(g)}}{6.02 \times 10^{23} \text{ part of CH}_3\text{OH}_{(l)} \quad 1 \text{ mol of CH}_3\text{OH}_{(l)}} = 1.3289 \dots \text{ mol} = 1.33 \text{ mol of H}_2\text{O}_{(g)}$$

STEP BY STEP:

2) $n = p/P = 4.00 \times 10^{23} \text{ part of CH}_3\text{OH}_{(l)} / 6.02 \times 10^{23} \text{ part of CH}_3\text{OH}_{(l)} \text{ per mol} = 0.66445 \dots \text{ mol of CH}_3\text{OH}_{(l)}$

3) $n_r = n_g \times R/G = 0.66445 \dots \text{ mol of CH}_3\text{OH}_{(l)} \times \frac{2 \text{ mol of H}_2\text{O}_{(g)}}{1 \text{ mol of CH}_3\text{OH}_{(l)}} = 1.3289 \dots \text{ mol} = 1.33 \text{ mol of H}_2\text{O}_{(g)}$

3. If 122.6 g of solid potassium chlorate is heated, the crystals melt and decompose into solid potassium chloride and oxygen gas. How many moles of potassium chloride are formed?



K = $39.10 \times 1 = 39.10$

Cl = $35.45 \times 1 = 35.45$

O = $16.00 \times 3 = 48.00$

LINEAR METHOD:

$$\frac{122.6 \text{ g of KClO}_{3(s)} \times 1 \text{ mol of KClO}_{3(s)} \times 2 \text{ mol of KCl}_{(s)}}{122.55 \text{ g of KClO}_{3(s)} \quad 2 \text{ mol of KClO}_{3(s)}} = 1.000 \text{ mol of KCl}_{(s)}$$


TOTAL 125.55g/mol

STEP BY STEP METHOD

2) $n = m/M = 122.6 \text{ g} / 122.55 \text{ g/mol} = 1.000 \text{ mol of KClO}_{3(s)}$

3) $n_R = n_G \times R/G = 1.000 \text{ mol of KClO}_{3(s)} \times \frac{2 \text{ mol of KCl}_{(s)}}{2 \text{ mol of KClO}_{3(s)}} = 1.000 \text{ mol of KCl}_{(s)}$

4. Black iron (III) oxide solid can be converted into water and iron metal when the iron (III) oxide is reacted with hydrogen gas. If 125 g of iron (III) oxide is reacted, how many moles of water are formed?



Fe = $55.85 \times 2 = 111.70$

O = $16.00 \times 3 = 48.00$

LINEAR METHOD:

$$\frac{125 \text{ g of Fe}_2\text{O}_{3(s)} \times 1 \text{ mol of Fe}_2\text{O}_{3(s)} \times 3 \text{ mol of H}_2\text{O}_{(g)}}{159.70 \text{ g of Fe}_2\text{O}_{3(s)} \quad 1 \text{ mol of Fe}_2\text{O}_{3(s)}} = 2.34815 \dots \text{ mol} = 2.35 \text{ mol of H}_2\text{O}_{(g)}$$


TOTAL 159.70g/mol

STEP BY STEP METHOD:

2) $n = m/M = 125 \text{ g} / 159.70 \text{ g/mol} = 0.7827175 \dots \text{ mol of Fe}_2\text{O}_{3(s)}$

3) $n_R = n_G \times R/G = 0.7827175 \dots \text{ mol of Fe}_2\text{O}_{3(s)} \times \frac{3 \text{ moles of H}_2\text{O}_{(g)}}{2 \text{ moles of Fe}_2\text{O}_{3(s)}} = 2.34815 \dots \text{ mol} = 2.35 \text{ mol of H}_2\text{O}_{(g)}$

5. How many moles of zinc can react with hydrochloric acid to form 44.8 L of hydrogen gas at STP?



LINEAR METHOD:


$$\frac{44.8 \text{ L of H}_2 \times 1 \text{ mol of H}_2}{22.4 \text{ L of H}_2} \times \frac{1 \text{ mol of Zn}}{1 \text{ mol of H}_2} = \mathbf{2.00 \text{ mol of Zn}}$$

STEP BY STEP METHOD:

2) $n = v/V = 44.8 \text{ L} / 22.4 \text{ L/mol} = 2.00 \text{ mol of H}_2$


3) $n_R = n_G \times R/G = \frac{2.00 \text{ mol of H}_2 \times 1 \text{ mol of Zn}}{1 \text{ mol of H}_2} = \mathbf{2.00 \text{ mol of Zn}}$

6. Solutions of copper (II) sulphate and potassium phosphate are mixed. If 8.5 g of copper (II) phosphate form, how many moles of copper (II) sulphate react?



LINEAR: $\frac{8.5 \text{ g of Cu}_3(\text{PO}_4)_{2(s)} \times 1 \text{ mol of Cu}_3(\text{PO}_4)_{2(s)}}{380.59\text{g/mol of Cu}_3(\text{PO}_4)_{2(s)}} \times \frac{1 \text{ mol of Sn}_3\text{N}_{2(s)}}{2 \text{ mol of Fe}_3\text{N}_{2(s)}} = \mathbf{0.250 \text{ mol of tin (IV) phosphate}}$

7. In the manufacturing of nitric acid, nitrogen dioxide gas reacts with water to form nitric acid and nitrogen monoxide gas. How many moles of nitrogen dioxide gas reacts if 120.6 L of nitrogen monoxide gas is formed at SATP?



LINEAR METHOD:


$$\frac{120.6 \text{ L of NO}_{(g)} \times 1 \text{ mol of NO}_{(g)}}{24.8 \text{ L of NO}_{(g)}} \times \frac{3 \text{ mol of NO}_{2(g)}}{1 \text{ mol of NO}_{2(g)}} = \mathbf{14.59 \text{ moles of NO}_{(g)}}$$

STEP BY STEP METHOD:

2) $n = v/V = 120.6 \text{ L} / 24.8 \text{ L/mol} = 4.863 \text{ mol of NO}_{(g)}$

3) $n_R = n_G \times R/G = \frac{4.863 \text{ mol of NO}_{(g)} \times 3 \text{ mol of NO}_{2(g)}}{1 \text{ mol of NO}_{2(g)}} = \mathbf{14.59 \text{ moles of NO}_{(g)}}$

8. The thermite reaction is used in welding iron and steel. Aluminium and iron (III) oxide are ignited at high temperatures to produce aluminium oxide and iron. If 15.0 g of aluminium is used in this reaction, how many moles of aluminium oxide will be produced?



Step 2) $n = m/M = 15.0\text{g of Al}_{(s)} / 26.98\text{g/mol} = 0.55596\dots\text{mol of Al}_{(s)}$


Step 3) $n_R = n_G \times R/G = \frac{0.55596 \text{ mol of Al}_{(s)} \times 1 \text{ mol of Al}_2\text{O}_{3(s)}}{2 \text{ mol of Al}_{(s)}} = \mathbf{0.27798\dots}$
 $= \mathbf{0.278 \text{ mol of Al}_2\text{O}_{3(s)}}$

LINEAR: $\frac{15.0\text{g of Al}_{(s)} \times 1 \text{ mol of Al}_{(s)}}{26.98\text{g of Al}_{(s)}} \times \frac{1 \text{ mol of Al}_2\text{O}_{3(s)}}{2 \text{ mol of Al}_{(s)}} = 0.27798\dots \text{ mol} = \mathbf{0.278 \text{ mol of Al}_2\text{O}_{3(s)}}$

Worksheet 2.4: Quantity to Quantity Stoichiometry

Directions: Solve the following hypothetical stoichiometry problems. Assume water is available.

1. How many particles of aluminium oxide must be decomposed to produce 80.0 g of oxygen gas at STP?



LINEAR METHOD:

$$\frac{80.0 \text{ g of O}_2 (\text{g}) \times 1 \text{ mol of O}_2 (\text{g}) \times 2 \text{ mol of Al}_2\text{O}_3 (\text{s}) \times 6.02 \text{ E} 23 \text{ particles of Al}_2\text{O}_3 (\text{s})}{32.00 \text{ g of O}_2 (\text{g}) \quad 3 \text{ mol of O}_2 (\text{g}) \quad 1 \text{ mol of Al}_2\text{O}_3 (\text{s})} = 1.00 \text{ E} 24 \text{ part of Al}_2\text{O}_3 (\text{s})$$


STEP BY STEP:

2) $n = m/M = 80.0 \text{ g} / 32.00 \text{ g/mol} = 2.5 \text{ mol of O}_2$

3) $n_R = n_G \times R/G = 2.5 \text{ mol of O}_2 \times \frac{2 \text{ mol of Al}_2\text{O}_3}{3 \text{ mol of O}_2} = 1.66666... \text{ mol of Al}_2\text{O}_3 (\text{s})$

4) $p = n P = 1.6666... \text{ mol} \times 6.02 \text{ E} 23 \text{ particles/mol} = 1.00 \text{ E} 24 \text{ or } 1.00 \times 10^{24} \text{ particles of Al}_2\text{O}_3 (\text{s})$

2. Natural gas is mainly made up of methane. What mass of methane must be burned to produce 56.0 L of carbon dioxide at STP?



LINEAR: $\frac{56.0 \text{ L of CO}_2 \times 1 \text{ mol of CO}_2 \times 1 \text{ mol of CH}_4 \times 16.05 \text{ g of CH}_4}{22.4 \text{ L of CO}_2 \quad 1 \text{ mol of CO}_2 \quad 1 \text{ mol of CH}_4} = 40.1 \text{ g of CH}_4 (\text{g})$


STEP BY STEP:

2) $n = v/V = 56.0 \text{ L of CO}_2 / 22.4 \text{ L of CO}_2 = 2.5 \text{ mol of CO}_2$

3) $n_R = n_G \times R/G = 2.5 \text{ mol of CO}_2 \times 1 \text{ mol of CH}_4 / 1 \text{ mol of CO}_2 = 2.5 \text{ mol of CO}_2$

4) $m = nM = 2.5 \text{ mol of CO}_2 \times 16.05 \text{ g/mol of CO}_2 = 40.1 \text{ g of CH}_4 (\text{g})$

3. Aluminium metal is refined from bauxite ore. In the refining process, aluminium oxide decomposes to aluminium and oxygen gas. What mass of aluminium can be produced from 2.04 kg of aluminium oxide?



LINEAR: $\frac{2040 \text{ g of Al}_2\text{O}_3 \times 1 \text{ mol of Al}_2\text{O}_3 \times 4 \text{ mol of Al} \times 26.98 \text{ g of Al}}{101.96 \text{ g of Al}_2\text{O}_3 \quad 2 \text{ mol of Al}_2\text{O}_3 \quad 1 \text{ mol of Al}} = 1079.6 \text{ g} = 1.08 \text{ kg of Al}$


STEP BY STEP:

2) $n = m/M = 2040 \text{ g of Al}_2\text{O}_3 / 101.96 \text{ g/mol of Al}_2\text{O}_3 = 20.0078... \text{ mol of Al}_2\text{O}_3$

3) $n_R = n_G \times R/G = 20.0078... \text{ mol of Al}_2\text{O}_3 \times 4 \text{ mol of Al} / 2 \text{ mol of Al}_2\text{O}_3 = 40.0156... \text{ mol of Al} (\text{s})$

4) $m = nM = 40.0156... \text{ mol of Al} (\text{s}) \times 26.98 \text{ g/mol of Al} = 1079.6 \text{ g} = 1.08 \text{ kg of Al}$

4. Sodium hydrogen carbonate can be used to neutralize acids. If sodium hydrogen carbonate reacts with hydrochloric acid, what volume of carbon dioxide gas at STP can be produced by 16.8 g of sodium hydrogen carbonate? NOTE: Sodium chloride and water vapour is also produced.



LINEAR: $\frac{16.8 \text{ g of NaHCO}_3 \times 1 \text{ mol of NaHCO}_3 \times 1 \text{ mol of CO}_2 \times 22.4 \text{ L of CO}_2}{84.01 \text{ g of NaHCO}_3 \quad 1 \text{ mol of NaHCO}_3 \quad 1 \text{ mol of CO}_2} = 4.48 \text{ L of CO}_2 (\text{g})$


STEP BY STEP:

2) $n = m/M = 16.8 \text{ g of NaHCO}_3 / 84.01 \text{ g/mol of NaHCO}_3 = 0.19997... \text{ mol of NaHCO}_3$

3) $n_R = n_G \times R/G = 0.19997... \text{ mol of NaHCO}_3 \times 1 \text{ mol of CO}_2 / 1 \text{ mol of NaHCO}_3 = 0.19997... \text{ mol of NaHCO}_3$

4) $v = nV = 0.19997... \text{ mol of NaHCO}_3 \times 22.4 \text{ L of CO}_2 = 4.48 \text{ L of CO}_2 (\text{g})$

5. Photography film is coated with silver chloride, which is produced when silver nitrate reacts with sodium chloride. What mass of silver chloride can be made from 11.7 g of sodium chloride?



$$\text{LINEAR: } \frac{11.7 \text{ g of NaCl} \times 1 \text{ mol of NaCl} \times 1 \text{ mol of AgCl} \times 143.32 \text{ g of AgCl}}{58.44 \text{ g of NaCl} \quad 1 \text{ mol of NaCl} \quad 1 \text{ mol of AgCl}} = 28.7 \text{ g of AgCl}$$


STEP BY STEP:

$$2) n = m/M = 11.7 \text{ g} / 58.44 \text{ g/mol} = 0.200205 \dots \text{ mol of NaCl}$$

$$3) n_R = n_G \times R/G = 0.200205 \dots \text{ mol of NaCl} \times \frac{1 \text{ mol of AgCl}}{1 \text{ mol of NaCl}} = 0.200205 \dots \text{ mol of AgCl}$$

$$4) m = nM = 0.200205 \dots \text{ mol of AgCl} \times 143.32 \text{ g/mol} = 28.693 \dots \text{ g} = 28.7 \text{ g of AgCl}$$

6. Ammonia reacts with hydrochloric acid to produce ammonium chloride. What volume of ammonia at SATP is needed to produce 36.1 g of ammonium chloride?



$$\text{LINEAR: } \frac{36.1 \text{ g of NH}_4\text{Cl} \times 1 \text{ mol of NH}_4\text{Cl} \times 1 \text{ mol of NH}_3 \times 24.8 \text{ L of NH}_3}{53.50 \text{ g of NH}_4\text{Cl} \quad 1 \text{ mol of NH}_4\text{Cl} \quad 1 \text{ mol of NH}_3} = 16.7 \text{ L of NH}_3$$


STEP BY STEP:

$$2) n = m/M = 36.1 \text{ g} / 53.50 \text{ g/mol} = 0.674766 \dots \text{ mol of NH}_4\text{Cl}$$

$$3) n_R = n_G \times R/G = 0.674766 \dots \text{ mol of NH}_4\text{Cl} \times \frac{1 \text{ mol of NH}_3}{1 \text{ mol of NH}_4\text{Cl}} = 0.674766 \dots \text{ mol of NH}_3$$

$$4) v = n V = 0.674766 \dots \text{ mol of NH}_3 \times 24.8 \text{ L/mol} = 16.734 \dots \text{ L} = 16.7 \text{ L of NH}_3$$

7. If sulphuric acid reacts with 29.4 g of potassium hydroxide, what mass of potassium sulphate is produced?



$$\text{LINEAR: } \frac{29.4 \text{ g of KOH} \times 1 \text{ mol of KOH} \times 1 \text{ mol of K}_2\text{SO}_4 \times 174.27 \text{ g of K}_2\text{SO}_4}{56.11 \text{ g of KOH} \quad 2 \text{ mol of KOH} \quad 1 \text{ mol of K}_2\text{SO}_4} = 45.7 \text{ g of K}_2\text{SO}_4$$


STEP BY STEP:

$$2) n = m/M = 29.4 \text{ g} / 56.11 \text{ g/mol} =$$

$$3) n_R = n_G \times R/G =$$

$$4) m = nM$$

8. If sodium iodide reacts with lead (II) nitrate, what mass of lead (II) nitrate will be required to produce 150 g of precipitate?



$$\text{LINEAR: } \frac{150 \text{ g of PbI}_2 \times 1 \text{ mol of PbI}_2 \times 1 \text{ mol of Pb(NO}_3)_2 \times 331.22 \text{ g of Pb(NO}_3)_2}{461 \text{ g of PbI}_2 \quad 1 \text{ mol of PbI}_2 \quad 1 \text{ mol of Pb(NO}_3)_2} = 108 \text{ g of Pb(NO}_3)_2$$

STEP BY STEP:

$$2) n = m/M =$$


$$3) n_R = n_G \times R/G =$$

$$4) m = nM$$

Worksheet 2.5: Limiting & Excess Reagents

Directions: For each of the following, write a balanced equation and determine the limiting reagent & the excess reagent (if they are present).

1. 5.0 mol of gasoline ($C_8H_{18(l)}$) burns 47.0 mol of oxygen at STP. How many moles of carbon dioxide are present at STP? **G** **G** **R**




$$\text{Step 3) } \frac{5.0 \text{ mol of } C_8H_{18} \times 8 \text{ mol of } CO_2}{1 \text{ mol of } C_8H_{18(l)}} = 40 \text{ mol of } CO_{2(g)}$$

$$\frac{47.0 \text{ mol} \times 8 \text{ mol of } CO_{2(g)}}{12.5 \text{ mol of } O_{2(g)}} = 30.08 \text{ mol of } CO_{2(g)} \text{ (LIMITING)} = \mathbf{30.1 \text{ mol of } CO_{2(g)}}$$

Step 4) not necessary.

2. 18.0 g of water breaks up into 6.0 g of oxygen. How many grams of hydrogen are formed?



$$\text{Step 2) } n = m/M = 18.0g/18.02g/mol \text{ of water} = 1.0...mol \text{ of water}$$

THIS IS NOT A LIMITING QUESTION BECAUSE THERE IS ONLY ONE REACTANT.

$$n = m/M = 6.0g/36.0g/mol = 0.166... \text{ mol of oxygen}$$

THIS QUESTION CAN NOT BE SOLVED BECAUSE YOU DON'T KNOW WHICH ONE IS GIVEN. THIS COULD BE A PERCENT YIELD QUESTION (NEXT LESSON.)

3. 22.4 mL of methane reacts with 22.4 mL of oxygen at SATP. How many moles of water are made? **G** **G** **R**



$$\text{Step 2) } \begin{array}{ll} n = v/V & n = v/V \\ = 0.0224/24.8 & = 0.0224/24.8 \\ = 9.03...E-4 \text{ mol} & = 9.03...E-4 \text{ mol} \end{array}$$


$$\text{Step 3) } \frac{9.03...E-4 \times 2 \text{ mol } H_2O_{(g)}}{1 \text{ mol } CH_{4(g)}} \quad \frac{9.03...E-4 \times 2 \text{ mol } H_2O_{(g)}}{2 \text{ mol } O_{2(g)}} \\ = 1.806..E-3 \text{ mol (EXCESS)} = \mathbf{9.03E-4 \text{ mol (LIMIT)}}$$

Step 4) If we changed it to volume


$$9.03...E-4 \text{ mol} \times 24.8 \text{ L/mol} = 0.0224 \text{ L of water or } 22.4 \text{ mL of water}$$

4. 26 g of magnesium react with 1.00 mol of hydrochloric acid.

a) What volume of Hydrogen gas is made at STP?




b) How much excess reagent is left over?



Step 4) Excess = Original – used = $1.0695... \text{ mol} - 0.500 \text{ mol} = 0.570 \text{ mol of Mg}_{(s)}$ left
 $m = n M = 0.5695... \text{ mol of Mg} \times 24.31 \text{ g/mol} = 13.8 \text{ g of Mg}_{(s)}$ left over

5. 3.02×10^{23} formula units of sodium react with 12 L of chlorine gas at STP. How much excess reagent is left over if the limiting is all used up?



Step 4) Excess = original – used = $1.0714... \text{ mol} - 0.2508... \text{ mol} = 0.821 \text{ mol of Cl}_{2(g)}$ left
 $v = nV$; $v = 0.821... \text{ mol of Cl}_{2(g)} \times 22.4 \text{ L/mol} = 18.380... \text{ L} = 18.4 \text{ L of Cl}_{2(g)}$


6. Describe a limiting reagent and an excess reagent.

- A limiting reagent is a reactant that controls how much product you have (it is the first reagent to be used up).
- An excess reagent is a reactant that is left over (it is not all used up)


Worksheet 2.6: Percent yield and Percent error

Directions: For each of the following write a balanced equation and determine the theoretical yield, actual yield, percent yield & the percent error.


1. 8.0 mol of sulphur dioxide decomposes and actually produces 7.0 mol of oxygen gas.




2. 26.0 g of aluminum reacts with a solution of calcium nitrate and produces 3.00 moles of calcium.




3. 6.50 mol of potassium chlorate solid is heated and breaks down into potassium chloride solid and 223 L of oxygen gas at SATP.




4. 33.6 L of methane burns and produces 2.00 mol of carbon dioxide gas at STP.



5. Sulphuric acid reacts with 29.4 g of potassium hydroxide and produces 40.5 g of potassium sulphate



- Step 4) Don't have to do this step because both AY and TY are in moles.



6. Describe percent yield and percent error.

Percent yield: a ratio between AY and TY as a percent; how much you produce compared to what you should produce.

Percent error: an indication of error (human, instrumental & experimental).

Worksheet 2.7: Limiting Reagents and Percent Yield

1. Methane gas burns at STP.
- a. If 0.500 mol of methane is burned in 2.50 mol of oxygen, what is the limiting reagent? **G1** **G2** **R**
- Step 1)** $1 \text{ CH}_{4(g)} + 2 \text{ O}_{2(g)} \rightarrow \text{CO}_{2(g)} + 2 \text{ H}_2\text{O}_{(g)}$
- Step 2)** $0.500 \text{ mol} \quad 2.50 \text{ mol}$
- Step 3)** $\frac{0.500 \times 2 \text{ mol of H}_2\text{O}_{(g)}}{1 \text{ mol of CH}_{4(g)}} = 1.00 \text{ mol of water}$ $\frac{2.50 \times 2 \text{ mol of H}_2\text{O}_{(g)}}{2 \text{ mol of O}_{2(g)}} = 2.50 \text{ mol of water}$
- CH₄ is limiting** **O₂ is excess**
- b. What is the theoretical yield, in moles, of water?
1.00 mol of water is the theoretical yield (use the limiting side)

2. Sodium and chlorine are mixed together.
- a. What is the limiting reagent if there is 10.0 g of sodium and 20.0 g of chlorine?
- G1** **G2** **R**
- 1) $2 \text{ Na}_{(s)} + 1 \text{ Cl}_{2(g)} \rightarrow 2 \text{ NaCl}_{(aq)}$
- 2) $n = m/M = 10.0\text{g}/22.99\text{g/mol} = 0.4349717... \text{ mol}$ $n = 20.0\text{g}/70.90\text{g/mol} = 0.282087447... \text{ mol}$
- 3) $\times 2 \text{ mol of NaCl}_{(aq)}/2 \text{ mol of Na}_{(s)} = 0.4349... \text{ mol of NaCl}_{(aq)}$ (smaller) $\times 2 \text{ mol of NaCl}_{(aq)}/1 \text{ mol of Cl}_{2(g)} = 0.56417... \text{ mol of NaCl}_{(aq)}$
- Na_s is LIMITING** **Cl_{2(g)} is EXCESS**

LINE METHOD:

$$10.0\text{g} \times \frac{1 \text{ mol of Na}_{(s)}}{22.99 \text{ g}} \times \frac{2 \text{ mol of NaCl}_{(aq)}}{2 \text{ mol of Na}_{(s)}} \times \frac{58.44 \text{ g of NaCl}_{(aq)}}{1 \text{ mol of NaCl}_{(aq)}} = 25.419... \text{ g of NaCl LIMIT}$$

$$20.0\text{g} \times \frac{1 \text{ mol of Cl}_2}{70.90 \text{ g}} \times \frac{2 \text{ mol of NaCl}_{(aq)}}{1 \text{ mol of Cl}_{2(s)}} \times \frac{58.44 \text{ g of NaCl}_{(aq)}}{1 \text{ mol of NaCl}_{(aq)}} = 32.970... \text{ g of NaCl EXCESS}$$

- b. How many grams of the product are produced?
- 4) $m = nM_{\text{NaCl}} = 0.4349717... \text{ mol of NaCl}_{(aq)} \times 58.44\text{g/mol} = 25.4 \text{ g of NaCl}_{(aq)}$

3. In the synthesis of sulphuric acid, one step involves the mixing of sulphur dioxide with oxygen to produce sulphur trioxide. If 175 L of sulphur dioxide was mixed with 85 L of oxygen at SATP, how many litres of sulphur trioxide is produced?

G1 = E **G2 = L** **R**

STEP 1) $2\text{SO}_{2(g)} + 1\text{O}_{2(g)} \rightarrow 2\text{SO}_{3(g)}$

$175 \text{ L} \quad 85\text{L} \quad ?$

STEP 2) $n = v/V = 175\text{L}/24.8\text{L/mol} = 7.0564... \text{ mol SO}_{2(g)}$ $n = 85\text{L}/24.8\text{L/mol} = 3.4274... \text{ mol of O}_{2(g)}$

STEP 3) $\frac{\times 2 \text{ mol SO}_{3(g)}}{2 \text{ mol SO}_{2(g)}} = 7.0564... \text{ mol}$ $\frac{\times 2 \text{ mol SO}_{3(g)}}{1 \text{ mol of O}_{2(g)}} = 6.8548... \text{ mol of SO}_{3(g)}$ (LIMITING)

STEP 4) $v = nV = 6.8548... \text{ mol of SO}_{3(g)} \times 24.8 = 170 \text{ L of SO}_{3(g)}$


LINEAR METHOD

$$\text{For SO}_{2(g)}: \frac{175\text{L} \times 1 \text{ mol of SO}_{2(g)} \times 2 \text{ mol of SO}_{3(g)} \times 24.8 \text{ L of SO}_{3(g)}}{24.8 \text{ L of SO}_{2(g)} \times 2 \text{ mol of SO}_{2(g)} \times 1 \text{ mol of SO}_{3(g)}} = 174.99 \dots \text{ L of SO}_{3(g)}$$

$$\text{For O}_{2(g)}: \frac{85\text{L} \times 1 \text{ mol of O}_{2(g)} \times 2 \text{ mol of SO}_{3(g)} \times 24.8 \text{ L of SO}_{3(g)}}{24.8 \text{ L of O}_{2(g)} \times 1 \text{ mol of O}_{2(g)} \times 1 \text{ mol of SO}_{3(g)}} = 170 \text{ L of SO}_{3(g)}$$


4. Adipic acid ($C_6H_{10}O_4(s)$), a raw material for nylon, is made by the oxidation (reacting with oxygen) of cyclohexane ($C_6H_{12(s)}$). Water is a by-product.

a. How many moles of oxygen gas would be needed to make 40.0 mol of adipic acid?






STEP 4) Not needed



LINEAR METHOD



b. If 2.00 mol of oxygen is reacted with 164 g of cyclohexane, what is the theoretical yield of adipic acid in grams? $G1 = L$



LINEAR METHOD




c. If 285 g of acid was produced in b) what is the percent yield?

(Use rounded answer from b)

$\% \text{ yield} = \text{AY/TY} \times 100\% = 285 \text{ g} / 1 \times 10^2 \text{ g of } C_6H_{10}O_4(s) \times 100\% = 285\% \text{ yield} = 3 \times 10^2 \%$


5. A chemist, new to the behaviour of chlorine toward hydrocarbon compounds, tried to make dichloromethane ($CH_2Cl_{2(g)}$), by mixing 5500 mL of chloromethane ($CH_3Cl_{(g)}$) and 5500 mL of chlorine at STP. Hydrogen chloride gas was a by product. After the reaction was complete, some chloromethane remained unchanged and 12.8 g of dichloromethane was obtained.

a. Which reactant is excess?



Chloromethane is excess

b. How much dichloromethane can theoretically be produced?



c. What is the percent yield?


$\% \text{ yield} = \text{AY/TY} \times 100\% = 12.8 \text{ g} / 13.9 \text{ g} \times 100\% = 92.086... \% = 92.1\%$

d. What is the percent error?


$\% \text{ error} = (\text{TY}-\text{AY})/\text{TY} \times 100\% = (13.9 - 12.8)/13.9 \times 100\% = 7.91\%$

Worksheet 3.8: Stoichiometry Review



1. A 34.5 g sample of sodium hydroxide solution is reacted with excess phosphoric acid. What is the mass of sodium phosphate that will form?





LINE METHOD:



2. A 75.0 g sample of lithium carbonate reacts with 120 g of aluminum nitrate. What mass of precipitate will form?




b) HOW MUCH EXCESS LEFT OVER





$$75.0 - 62.4 = 12.6 \text{ g of EXCESS LEFT OVER}$$


3. A 45.0 g sample of ethanol burns in the presence of 105 L of oxygen gas. (assume STP). What is the percent yield if 50.0 g of carbon dioxide is formed?



LINE METHOD:



4. When 5.6×10^{24} particles of magnesium sulfide reacts with potassium hydroxide, then 500 g of precipitate forms. What is the percent error?



$$\text{Step 2) } n=p/P=5.6E24/6.02E23 \quad \text{500g = AY}$$

$$n = 9.3023... \text{ mol}$$

$$\text{Step 3) R/G } \begin{array}{l} \times 1 \text{ mol of Mg(OH)}_{2(s)} = 9.3023... \text{ mol of Mg(OH)}_{2(s)} \\ 1 \text{ mol of MgS} \end{array}$$


$$\text{Step 4) } m=nM; m = 9.23023... \text{ mol of Mg(OH)}_{2(s)} \times 58.33 \text{ g/mol of Mg(OH)}_{2(s)} = 542.604... \text{ g} = \text{TY}$$

LINE METHOD:

$$\frac{5.6E24 \text{ of MgS} \times 1 \text{ mol of MgS} \times 1 \text{ mol of Mg(OH)}_{2(s)} \times 58.33 \text{ g of Mg(OH)}_{2(s)}}{6.02E23 \text{ of MgS} \quad 1 \text{ mol of MgS} \quad 1 \text{ mol of Mg(OH)}_{2(s)}} = 542.604... \text{ g of Mg(OH)}_{2(s)}$$

$$\text{Step 5) } \% \text{error} = (\text{TY}-\text{AY})/\text{TY} \times 100\% = (542.604... \text{ g} - 500\text{g})/542.604... \text{ g} \times 100\% = 7.85\%$$

5. When 36.9 L of chlorine gas (SATP) reacts with 36.8 g of magnesium oxide, 38.9 g of magnesium chloride formed. What mass of magnesium chloride did you expect?



$$\text{Step 2) } n=v/V; n=36.9\text{L}/24.8\text{L/mol} \quad n=m/M; n=36.8\text{g}/40.31\text{g/mol} \quad 38.9 \text{ g} = \text{AY}$$

$$n=1.4879... \text{ mol of Cl}_{2(g)} \quad n=0.9129... \text{ mol of MgO}$$

$$\text{Step 3) R/G } 0.9129... \text{ mol of MgO} \times 2 \text{ mol of MgCl}_{2(s)} / 2 \text{ mol of MgO} = 0.9129... \text{ mol of MgCl}_{2(s)}$$


$$\text{Step 4) } m=nM; m = 0.9129... \text{ mol of MgCl}_{2(s)} \times 95.21 \text{ g/mol} = 86.9195... \text{ g} = 86.9 \text{ g of MgCl}_{2(s)} = \text{TY}$$

LINE METHOD:

$$\text{For Cl}_2: \frac{36.9\text{L of Cl}_2 \times 1 \text{ mol of Cl}_{2(g)} \times 2 \text{ mol of MgCl}_{2(s)} \times 95.21 \text{ g of MgCl}_{2(s)}}{24.8\text{L of Cl}_{2(g)} \quad 2 \text{ mol of Cl}_{2(g)} \quad 1 \text{ mol of MgCl}_{2(s)}} = 141.66 \text{ g of MgCl}_{2(s)}$$

$$\text{For MgO: } \frac{36.8\text{g of MgO} \times 1 \text{ mol of MgO} \times 2 \text{ mol of MgCl}_{2(s)} \times 95.21 \text{ g of MgCl}_{2(s)}}{40.31\text{g of MgO} \quad 2 \text{ mol of MgO} \quad 1 \text{ mol of MgCl}_{2(s)}} = 86.9 \text{ g of MgCl}_{2(s)}$$

6. When 24.5 g of iron(II) chloride reacts with 35.0 g of zinc, 7.2 g of iron was formed. What is the percent yield and percent error is this experiment?



$$\text{Step 2) } n=m/M; n = 24.5\text{g}/126.75\text{g/mol} \quad n=35.0\text{g}/65.41\text{g/mol} \quad 7.2\text{g} = \text{AY}$$

$$n=0.19329... \text{ mol} \quad n = 0.53508... \text{ mol}$$

$$\text{Step 3) R/G } 0.19329... \text{ mol} \times 1 \text{ mol of Fe}_{(s)} / 1 \text{ mol of FeCl}_{2(aq)} = 0.19329... \text{ mol of Fe}$$

$$\text{Step 4) } m=nM; m=0.19329... \text{ mol of Fe} \times 55.85 \text{ g/mol} = 10.79... \text{ g of Fe} = \text{TY}$$

LINE METHOD:

$$\text{For FeCl}_2: \frac{24.5 \text{ g of FeCl}_{2(aq)} \times 1 \text{ mol of FeCl}_{2(aq)} \times 1 \text{ mol of Fe}_{(s)} \times 55.85 \text{ g of Fe}_{(s)}}{126.75 \text{ g of FeCl}_{2(aq)} \quad 1 \text{ mol of FeCl}_{2(aq)} \quad 1 \text{ mol of Fe}_{(s)}} = 10.79... \text{ g of Fe}_{(s)} = \text{TY}$$

$$\text{For Zn: } \frac{35.0\text{g of Zn}_{(s)} \times 1 \text{ mol of Zn}_{(s)} \times 1 \text{ mol of Fe}_{(s)} \times 55.85\text{g of Fe}_{(s)}}{65.41\text{g of Zn}_{(s)} \quad 1 \text{ mol of Zn}_{(s)} \quad 1 \text{ mol of Fe}_{(s)}} = 29.8845... \text{ of Fe}_{(s)}$$

$$\text{Step 5) } \% \text{yield} = \text{AY}/\text{TY} \times 100\% = 7.2\text{g}/10.79... \text{ g} \times 100\% = 66.69... \% = 66.7\%$$

$$\% \text{error} = 100 - \% \text{yield} = 33.3\%$$

Worksheet 2.11: Introduction to Gases & Dalton's Gas Law

1. What are three physical properties of all gases?
Gases do not have a fixed volume or shape (fill container), are compressible and diffuse.

2. What three variables affect gases?
Pressure, temperature and volume

3. What instrument measures pressure?
Barometer (manometer)

4. What is the SI unit for pressure?
KPa (kiloPascals)

5. What unit expresses the average kinetic energy of a gas?
Kelvin or degrees Celcius

6. A 1.00 L bottle of gas contains oxygen at 10.0 kPa, nitrogen at 12.1 kPa and hydrogen at 97.5 kPa
a. What is the total pressure?

$$P_T = P_1 + P_2 + P_3$$

$$P_T = 10.0 \text{ kPa} + 12.1 \text{ kPa} + 97.5 \text{ kPa}$$

$$P_T = 119.6 \text{ kPa (one decimal place for significant digits)}$$

- b. What percent of each gas is present? (HINT % = $P_{\text{gas}}/P_{\text{total}} \times 100$)

$$\text{oxygen} = 8.36 \%; \text{ nitrogen} = 10.1 \%; \text{ hydrogen} = 81.5 \% \text{ (3 significant digits)}$$

- c. What is the volume of each gas?

$$V \text{ of oxygen} = 1.00\text{L} \times 0.0836 = 0.0836 \text{ L} = \underline{83.6 \text{ mL}}$$

7. Four gases (A,B,C and D) make up a mixture with a pressure of 150 kPa. What is the partial pressure of gas A, if gas B has a pressure of 58.0 kPa, gas C has a pressure of 23.8 kPa and gas D has a pressure of 15.9 kPa.

$$P_1 = P_T - (P_2 + P_3 + P_4)$$

$$P = 150\text{kPa} - (58.0\text{kPa} + 23.8\text{kPa} + 15.9\text{kPa})$$

$$P = \underline{52.3 \text{ kPa}}$$

8. Three gases make up a mixture. At a particular pressure, the partial pressures are measured: Gas A = 67.00 kPa, Gas B, 6.70 kPa, and Gas C = 0.67 kPa. What is the pressure conditions under which this measurement is taken?

$$P_T = P_1 + P_2 + P_3$$

$$P = 67.00 \text{ kPa} + 6.70 \text{ kPa} + 0.67 \text{ kPa} = \underline{74.37 \text{ kPa}}$$

Worksheet 2.12: Boyles' Gas Law

1. What is the pressure when: (temperature is constant)

a. 130 mL of a gas at 740 mmHg is changed to 150 mL?

$$P_2 = P_1V_1/V_2; P_2=740\text{mmHg} \times 130\text{mL} / 150\text{mL}$$

$$P_2=\underline{641 \text{ mmHg}}$$

b. 25 mL of gas at 65 atm is changed to 30.0 mL?

$$P_2 = P_1V_1/V_2; P_2=65\text{atm} \times 25\text{mL} / 30.0\text{mL}$$

$$P_2=\underline{54 \text{ atm}}$$

c. 1.0 L of gas at 70 kPa is changed to 1.2 L?

$$P_2 = P_1V_1/V_2; P_2=70\text{kPa} \times 1.0\text{L} / 1.2\text{L}$$

$$P_2=\underline{58 \text{ kPa}}$$

2. What is the volume when: (temperature is constant)

a. 75 mL of gas at 4.1 atm is changed to 7.0 atm?

$$V_2 = P_1V_1/P_2; V_2=4.1\text{atm} \times 75\text{mL} / 7.0\text{atm}$$

$$V_2=\underline{44 \text{ mL}}$$

b. 60.0 mL of gas at 760 mmHg is changed to 10 mmHg?

$$V_2 = P_1V_1/P_2; V_2=760\text{mmHg} \times 60\text{mL} / 10 \text{ mmHg}$$

$$V_2=\underline{4.6\text{L} \text{ or } 4.6 \times 10^3 \text{ mL}}$$

c. 400.0 mL of gas at 760 kPa is changed to 300 kPa?

$$V_2 = P_1V_1/P_2; V_2=760\text{kPa} \times 400\text{ml}/300\text{kPa}$$

$$V_2=\underline{1013.33 \text{ mL}; 1.01 \times 10^3 \text{ mL} \text{ or } 1.01 \text{ L}}$$

Worksheet 2.13: Charles' Gas Law

1. What is the volume when: (pressure is constant)
- a. 125 mL of gas at 25° C is cooled to Standard temperature?
 $T_1=25 + 273.15=298.15K$ $T_2=0 + 273.15=273.15K$
 $V_2 = V_1T_2/T_1$; $V_2=$
 $V_2=0.11L$ or 1.1×10^2 mL
- b. 300.0 mL of gas at 0.0°C is heated to 30.0°C?
 $T_1=0 + 273.15=273.15K$ $T_2=30 + 273.15=303.15K$
 $V_2 = V_1T_2/T_1$; $V_2=$
 $V_2=0.333L$ or $333mL$
- c. 220.0 mL of gas at 10.0°C is heated to 100.0°C?
 $T_1=10 + 273.15=283.15K$ $T_2=100 + 273.15=373.15K$
 $V_2 = V_1T_2/T_1$; $V_2=$
 $V_2=0.290L$ or 290 mL
2. What is the temperature when: (pressure is constant)
- a. 30.0 mL (V1) of gas at 14°C (T1) is compressed to 22 mL (V2)?
 $T_1=14 + 273.15=287.15K$
 $T_2 = V_2T_1/V_1$; $T_2=22mL \times 287.15/30mL$
 $T_2=210.6$ K; 2.1×10^2K or $-63C$
- b. 16.4 mL of gas at 28°C is expanded to 20.0 mL?
 $T_1=28 + 273.15=301.15K$
 $T_2 = V_2T_1/V_1$; $T_2=20 \times 301.15 / 16.4$
 $T_2=367.25K$; 3.7×10^2K or $94C$
- c. 39 mL of gas at 0.0°C is compressed to 35 mL?
 $T_1=0 + 273.15=273.15K$
 $T_2 = V_2T_1/V_1$; $T_2=35 \times 273.15/39$
 $T_2=245.1K$; 2.5×10^2K or $-28C$

Worksheet 2.14: Lusac's Gas Law

1. What is the pressure when: (volume is constant)

a. a gas at 130 C and 740 mmHg is changed to 150 C?

$$T_1 = 273.15 + 130 = 403.15\text{K} \quad T_2 = 273.15 + 150 = 423.15\text{K}$$

$$P_1 = P_2 T_1 / T_2; \quad P_1 = 740\text{mmHg} \times 403.15\text{K} / 423.15\text{K}$$

$$P_1 = \underline{777 \text{ mmHg}}$$

b. a gas at 25 C and 65 atm is changed to 30.0 C?

$$T_1 = 273.15 + 25 = 293.15\text{K} \quad T_2 = 273.15 + 30.0 = 303.15\text{K}$$

$$P_1 = P_2 T_1 / T_2; \quad P_1 =$$

$$P_1 = \underline{66 \text{ atm}}$$

c. a gas at 1.0 K and 70 kPa is changed to 1.2 K?

$$P_1 = P_2 T_1 / T_2; \quad P_1 =$$

$$P_1 = \underline{84 \text{ kPa}}$$

2. What is the temperature in degrees Celcius when: (volume is constant)

a. a gas at 75.0 C and 4.10 atm is changed to 7.00 atm?

$$T_1 = 273.15 + 75.0 = 348.15\text{K}$$

$$T_2 = P_2 T_1 / P_1; \quad T_2 = 7.00 \times 348.15 / 4.10$$

$$T_2 = \underline{594 \text{ K or } 321\text{C}}$$

b. a gas at 60.0 C and 760 mmHg is changed to 10.0 mmHg?

$$T_1 = 273.15 + 60.0 = 333.15\text{K}$$

$$T_2 = P_2 T_1 / P_1; \quad T_2 =$$

$$T_2 = \underline{4.38\text{K or } -269 \text{ C}}$$

c. a gas at 113 K and 760 kPa is changed to 300 kPa?

$$T_2 = P_2 T_1 / P_1; \quad T_2 = 300\text{kPa} \times 113\text{K} / 760\text{kPa}$$

$$T_2 = \underline{44.6\text{K or } -229 \text{ C}}$$

Worksheet 2.15: Combined Gas Law

Solve the following gas problems.

1. If 120 mL of oxygen is collected at 27°C and 740 mmHg, what will the volume of the dry gas be at STP? **T=0C, P=760 mmHg**
 $T_1=273.15 + 27 = 300.15 \text{ K}$, $V_1=0.120\text{L}$, $P_1=740\text{mmHg}$
 $T_2=273.15 + 0 = 273.15 \text{ K}$, $V_2=?$, $P_2=760\text{mmHg}$
 $P_1V_1T_2=P_2V_2T_1$; $V_2=$
 $V_2=0.11\text{L}$ or $1.1 \times 10^2 \text{ mL}$
2. If 500.0 mL of hydrogen is collected at 293.15 K and 95.0 kPa, what will the volume of the gas be at SATP? **T=25C, P=100 kPa**
 $T_1=293.15 \text{ K}$, $V_1=0.500\text{L}$, $P_1=95.0\text{kPa}$
 $T_2=273.15 + 25 = 298.15 \text{ K}$, $V_2=?$, $P_2=100\text{kPa}$
 $P_1V_1T_2=P_2V_2T_1$; $V_2=$
 $V_2=0.483\text{L}$ or 483 mL
3. 113 mL of oxygen is collected at 22°C and 98.0 kPa and left over night. The next day, the volume was 109 mL and the temperature was 21°C. What was the pressure?
 $T_1=273.15 + 22 = 295.15 \text{ K}$, $V_1=0.113\text{L}$, $P_1=98.0\text{kPa}$
 $T_2=273.15 + 21 = 294.15 \text{ K}$, $V_2=0.109\text{L}$, $P_2=?$
 $P_1V_1T_2=P_2V_2T_1$; $P_2=$
 $P_2=101.25 \text{ kPa}$, $P_2=1.0 \times 10^2 \text{ kPa}$
5. 36 mL of nitrogen was collected at 25°C, but the barometer was broken so the pressure could not be read. Three days later, the new barometer arrived. The new volume was 32 mL, the temperature was 21°C and the pressure reading was 739 mmHg. What was the original pressure?
 $T_1=273.15 + 25 = 298.15 \text{ K}$, $V_1=0.036\text{L}$, $P_1=?$
 $T_2=273.15 + 21 = 294.15 \text{ K}$, $V_2=0.032\text{L}$, $P_2=739\text{mmHg}$
 $P_1V_1T_2=P_2V_2T_1$
 $P_1=665.8 \text{ mmHg}$, $P_1 = 6.7 \times 10^2 \text{ mmHg}$

4. If 250 mL of helium was collected at STP, what will the temperature be if the volume is reduced to 200 mL and the pressure increased to 110 kPa?

$$T_1 = 273.15 + 0 = 273.15 \text{ K}, V_1 = 0.250 \text{ L}, P_1 = 101.325 \text{ kPa}$$

$$T_2 = ?, V_2 = 0.200 \text{ L}, P_2 = 110 \text{ kPa}$$

$$P_1 V_1 T_2 = P_2 V_2 T_1; T_2 = 110 \text{ kPa} \times 0.200 \text{ L} \times 273.15 \text{ K} / (101.325 \text{ kPa} \times 0.250 \text{ L})$$

$$T_2 = \underline{237 \text{ K or } -35.9 \text{ C}}$$

5. A certain 1.0L sample of gas has a temperature of 23°C and a pressure of 0.96 atm. The sample was left overnight and the next day had a pressure of 1.00 atm and a volume of

1.1 L. What is the temperature on the second day?

$$T_1 = 273.15 + 23 = 296.15 \text{ K}, V_1 = 1.0 \text{ L}, P_1 = 0.96 \text{ atm}$$

$$T_2 = ?, V_2 = 1.1 \text{ L}, P_2 = 1.00 \text{ atm}$$

$$P_1 V_1 T_2 = P_2 V_2 T_1; T_2 = 1.1 \text{ L} \times 1.00 \text{ atm} \times 296.15 \text{ K} / (0.96 \text{ atm} \times 1.0 \text{ L})$$

$$T_2 = 339 \text{ K}; T_2 = \underline{3.4 \times 10^2 \text{ K or } 66 \text{ C}}$$

Worksheet 2.16: Ideal Gas Law

Solve the following gas problems.

1. What pressure (kPa) is exerted by 1.0 mol of an ideal gas contained in a 1.0 L vessel at 0.0°C?

$$PV=nRT; P=1.0\text{mol} \times 8.314\text{LkPa/Kmol} \times 273.15\text{K} / 1.0\text{L}$$

$$P=2270.969\text{ kPa}, \underline{P=2.3 \times 10^3\text{ kPa}}$$

2. What volume will 5.0 mol of an ideal gas occupy at 25.0°C and 1.5 atm of pressure?

$$V=nRT/P; V=5.0\text{mol} \times 0.0821\text{Latm/Kmol} \times 298.15\text{K} / 1.5\text{atm}$$

$$V=81.59\text{L}; \underline{V=82\text{L}}$$

3. Calculate the molar mass of gas if 4.5 L of the gas is at 785 mmHg, 23.5°C and the gas has a mass of 13.5 g.

$$n=PV/RT; n=785\text{mmHg} \times 4.5\text{L} / (62.4\text{LmmHg/Kmol} \times 296.65\text{K})$$

$$n=0.1915755\dots\text{mol}$$

$$M=m/n; M=13.5\text{g} / 0.19\dots\text{mol}; \underline{M=70.74\text{g/mol or }71\text{g/mol}}$$

4. 0.453 mol of a gas confined to a 15.0 L container exerts a pressure of 1.24 atm on the walls of the container. What is the temperature of the gas?

$$T=PV/nR; T=1.24\text{atm} \times 15.0\text{L} / (0.453\text{mol} \times 0.0821\text{Latm/Kmol})$$

$$\underline{T=500\text{K or }227\text{C}}$$

6. 5.4 g of carbon dioxide gas is confined to a 20.0 L container at a temperature of 315.5 K. What pressure (kPa) does the gas exert?

$$n=m/M; n=5.4\text{g}/44.01\text{g/mol}; n=0.122699\dots\text{mol}$$

$$P=nRT/V; P=0.12\dots\text{mol} \times 8.314\text{LkPa/Kmol} \times 315.15\text{K}/20.0\text{L}$$

$$P=16.09\text{ kPa}; \underline{P=16\text{ kPa}}$$

5. 2.125 g of a gas in a 1.25 L container exerts a pressure of 86.0 kPa at 40.0°C. What is the molar mass of the gas?

$$n=PV/RT; n=86.0\text{kPa} \times 1.25\text{L} / (8.314\text{LkPa/Kmol} \times 313.15\text{K})$$

$$n=0.041290\dots\text{mol}$$

$$M=m/n; M=2.125\text{g}/0.041290\dots\text{mol}; M=51.465\text{ g/mol}$$

$$M=\underline{51.46\text{g/mol or }51.5\text{ g/mol}}$$

6. To what temperature must 10.0 g of ammonia gas have to be heated in a 15.0 L container in order for it to exert a pressure of 3.50 atm?

$$n=m/M; n=10.0\text{g} / 17.04\text{g/mol}; n=0.5868\dots\text{mol}$$

$$T=PV/nR; T=3.50\text{atm} \times 15.0\text{L} / (0.5868\dots\text{mol} \times 0.0821\text{Latm/Kmol})$$

$$T=1089.6\text{ K}; \underline{T=1.09 \times 10^3\text{K}}$$

7. $2.0 \times 10^{-5}\text{g}$ of hydrogen gas at 327 K exerts a pressure of 50.5 kPa on the walls of a small tube. What is the volume of the tube?


$$n=m/M; n=m/M; n=2.0 \times 10^{-5}\text{g} / 2.02\text{g/mol}; n=9.9\dots \times 10^{-6}\text{ mol}$$

$$V=nRT/P; V=9.9\dots \times 10^{-6}\text{mol} \times 8.314\text{LkPa/Kmol} / 50.5\text{kPa}$$

$$\underline{V=5.3 \times 10^{-4}\text{ L}}$$

Worksheet 2.17: Gas Stoichiometry

1. What mass of propane from a tank can be burned using 50 L of oxygen at STP?



Step 2) ? $n = PV/RT$ or $n = v/V$

$n = 50\text{L}/22.4\text{L/mol}; n = 2.3\dots\text{mol}$

Step 3) $X/1\text{mol} = 2.3\dots\text{mol}/5\text{mol}$

$X = 0.44\dots\text{mol}$

Step 4) $m = nM; m = 0.44\dots\text{mol} \times 44.11\text{g/mol}$

$m = 19.69\text{ g}; m = 20\text{g}$

2. Hydrogen gas is burned in pollution-free vehicles to produce water vapor. What volume of hydrogen at 40°C and 150 kPa can be burned using 300 L of oxygen gas measured at the same conditions?



Step 2) ? $n = PV/RT$

$N = (150\text{kPa} \times 300\text{L}) / (8.314\text{LkPa/Kmol} \times 313.15\text{K})$

$N = 17.2\dots\text{mol}$

Step 3) $X/2\text{mol} = 17.2\dots\text{mol}/1\text{mol}$

$X = 34.5\dots\text{mol}$

Step 4) $V = nRT/P$

$V = 34.5\dots\text{mol} \times 8.314\text{LkPa/Kmol} \times 313.15\text{K} / 150\text{kPa}$

$V = 600\text{L}; V = 6.0 \times 10^2\text{L}$

3. A Down's Cell is used in the industrial production of sodium from the decomposition of molten sodium chloride. What is the temperature of 250 L of chlorine gas produced at 100.1 kPa if 100.0 g of sodium is also produced?



Step 2) $n = m/M$? K

$n = 100\text{g}/22.99\text{g/mol}$

$n = 4.3\dots\text{mol}$

Step 3) $4.3\dots\text{mol}/2\text{mol} = x/1\text{mol}$


$X = 2.1\dots\text{mol}$

Step 4) $T = PV/nR$

$T = 100.1\text{kPa} \times 250\text{L} / (2.1\dots\text{mol} \times 8.314)$

$T = 1.38 \times 10^3\text{ K or } 1.11 \times 10^3\text{ C}$

4. A typical home is heated with natural gas and consumes 2.00 ML of natural gas during the month of December. What volume of oxygen at STP is required to burn 2.00 ML of methane measured at 0°C and 120 kPa?



Step 2) $n = PV/RT$?
 $N = 120\text{kPa} \times 2.00 \times 10^6 / (8.314 \times 273.15\text{K})$
 $N = 105681 \dots \text{mol}$

Step 3) $105681 \dots \text{mol} / 1 \text{mol} = x / 2 \text{mol}$


$X = 211362 \dots \text{mol}$

Step 4) $V = nV$ or $V = nRT/P$

$V = 211362 \dots \text{mol} \times 22.4 \text{L/mol}$

$V = 4.73 \times 10^6 \text{ L}$ or 4.73 ML

5. Methane reacts with steam to produce hydrogen gas and carbon dioxide gas. What volume of hydrogen gas, measured at 25°C and 120 kPa, can be produced from 1.0 t of steam?



Step 2) $n = m/M$
 $n = 1000000\text{g} / 18.02\text{g/mol}$
 $n = 55493 \dots \text{mol}$

Step 3) $55493 \dots \text{mol} / 2 \text{mol} = x / 4 \text{mol}$
 $X = 11098 \dots \text{mol}$

Step 4) $V = nRT/P$
 $V = 11098 \times 8.314 \times 298.15\text{K} / 120\text{kPa}$
 $V = 2.3 \text{ ML}$ or $2.3 \times 10^6 \text{ L}$

6. Hydrogen gas can be produced from the electrolytic decomposition of water. What volume of hydrogen gas is produced, along with 52 kL of oxygen gas, at 25°C and 120kPa?



Step 2) ?
 $n = PV/RT$
 $n = (120\text{kPa} \times 52000\text{L}) / (8.314 \times 298.15)$
 $n = 2517 \dots \text{mol}$

Step 3) $X / 2 \text{mol} = 2517 \dots \text{mol} / 1 \text{mol}$
 $X = 5034 \dots \text{mol}$

Step 4) $V = nRT/P$; $V = 5034 \dots \text{mol} \times 8.314 \times 298.15\text{K} / 120\text{kPa}$
 $V = 104000\text{L}$; $V = 1.0 \times 10^5 \text{ L}$ or 0.10 ML

Worksheet 2.18: Review of Gases

1. A volume of 20.0 L of oxygen is warmed from -30.0 C to 85.0 C. What is the new volume, if the pressure is kept constant?

$$V_2 = V_1 T_2 / T_1; V_2 =$$

$$V_2 = 29.5 \text{ L}$$

2. A mass of air occupies a volume of 5.7 L at a pressure of 0.52 atm. What is the new pressure if the same mass of air at the same temperature is transferred to a 2.0 L container?

$$P_2 = P_1 V_1 / V_2; P_2 =$$

$$P_2 = 1.5 \text{ atm}$$

3. Determine the total pressure of a gas mixture that contains CO, Ne and He if the partial pressures of the gases are $P_{\text{CO}} = 1.53 \text{ atm}$, $P_{\text{Ne}} = 0.82 \text{ atm}$, and $P_{\text{He}} = 0.34 \text{ atm}$.

$$P_t = P_1 + P_2 + P_3; P_t =$$

$$P_t = 2.69 \text{ atm (2 decimal places are significant because you are adding.)}$$

4. What is the volume of a sample of oxygen gas that has a mass of 50.0 g and is under a pressure of 1.20 atm at 27.0 C?

$$V = nRT/P; V =$$

$$V = 32.1 \text{ L}$$

7. What is the volume at STP of a sample of carbon dioxide gas that has a volume of 75.0 mL at 30.0 C and 680 mmHg?

$$V_1 = P_2 V_2 T_1 / (T_2 P_1); V_1 =$$

$$V_1 = 0.0605 \text{ L or } 60.5 \text{ mL}$$

8. A rigid container holds a gas at a pressure of 0.55 atm at a temperature of -100 C. What will the pressure be when the temperature is increased to 200 C?

$$P_2 = P_1 T_2 / T_1; P_2 =$$

$$P_2 = 1.5 \text{ atm}$$


9. Explain why real gases deviate from the gas laws.

Real gases deviate because they can be condensed into liquids, have particle size and attract each other, unlike ideal gases.

Worksheet 4.1 – Solution Terminology and Theory

1. Illustrate (with a drawing) the difference between:

a) solute vs. solvent



b) homogenous mixture vs. heterogenous mixture

c) electrolyte vs. non-electrolyte

2. Illustrate two factors that affect the rate of solubility.

3. Illustrate how the following solids dissolve in water

a) Glucose

b) copper (II) sulphate

c) hydrochloric acid

4. Many reactions only occur when the reactants are dissolved in water. Why?

Worksheet 2.2: Concentration Problems

1. What is the molar concentration of an electroplating solution in which 1.50 mol of copper (II) sulphate are dissolved in 2.00 L of water?

$C=n/V$; $C=1.50\text{mol}/2.00\text{L}$; $C=0.750\text{ mol/L}$

2. What is the molar concentration of a solution in which 0.240 mol of washing soda, sodium carbonate decahydrate, is dissolved in 480 mL of water to make soft water solution?

$C=n/V$; $C=0.240\text{ mol}/0.480\text{ L}$; $C=0.500\text{ mol/L}$

3. What is the molar concentration of 500 mL of a solution that contains 12.7 g of swimming pool chlorinator, $\text{Ca}(\text{OCl})_2$?

1) $n=m/M$; $n=12.7/142.98\text{g/mol}$; $n=0.0888\dots\text{mol}$

Ca = 40.08

Ox2 = 32.00

Clx2 =70.90

2) $C=n/V$; $C=0.0888\dots\text{mol} / 0.500\text{L}$; $C=0.178\text{ mol/L}$

142.98

4. A given sample of household ammonia contains 156 g of ammonia dissolved in water to form a 2.00L solution. What is the molar concentration of the ammonia solution?

1) $n=m/M$; $n=156\text{g}/17.04\text{ g/mol}$; $n=9.154\dots\text{mol}$

N = 14.01

Hx3=3.03

17.04 g/mol

2) $C=n/V$; $C=9.154\dots\text{mol} /2.00\text{L}$; $C=4.58\text{ mol/L}$

5. Find the number of moles of sodium phosphate in 2.00L of a 0.100 mol/L sodium phosphate cleaning solution.
 $n = CV$; $n = 0.100 \text{ mol/L} \times 2.00 \text{ L}$; $n = 0.200 \text{ mol}$

6. How many moles of potassium sulphate are there in 500 mL of a 0.242 M solution used to remove rust stains?
 $n = CV$; $n = 0.242 \text{ mol/L} \times 0.500 \text{ L}$; $n = 0.121 \text{ mol}$
7. What mass of sodium bicarbonate must be added to a 2.50 L bowl to obtain a necessary 0.150 mol/L solution?
 $n = CV$; $n = 0.150 \text{ mol/L} \times 2.50 \text{ L}$; $n = 0.375 \text{ mol}$
 $m = nM$; $m = 0.375 \text{ mol} \times 84.01 \text{ g/mol} = 31.5 \text{ g}$
- | |
|-------------------------|
| Na = 22.99 |
| H = 1.01 |
| C = 12.01 |
| <u>Ox3=48.00</u> |
| 84.01 g/mol |
8. What volume of a 0.075 mol/L solution would contain the necessary 1.10 mol of sodium phosphate used to remove radiator scales?
 $V = n/C$; $V = 1.10 \text{ mol} / 0.075 \text{ mol/L}$; $V = 15 \text{ L}$
9. What mass of sodium silicate is necessary to prepare 10.0 L of a 0.00500 mol/L water softening solution?

10. How many litres of 0.800 mol/L solution would contain 119.2 g of NaOCl?

1) $n = m/M$; $n = 119.2\text{g}/74.44\text{g/mol}$; $n = 1.60\dots\text{mol}$

Na=22.99

O=16.00

2) $V = n/C$; $v = 1.60\dots\text{mol}/0.800\text{mol/L}$; $v = 2.00\text{ L}$

Cl=35.45

74.44g/mol

Worksheet 2.3: Making solutions and dilutions

1. A scientist has a container with solid sodium hydroxide and a container of 5.00 mol/L sodium hydroxide.
a) What are the two ways that the scientists can use to make a solution with a specific volume and concentration?

Make a solution by mixing a solute of specific mass with a specific volume of solvent (water)

OR make a dilution by adding water to a solution that is already made.

- b) What are two ways that the scientist can dilute the 5.00 mol/L solution?

Evaporate the solvent and then remove some solute and add the solvent back OR add more solvent to a small portion of the solution.

2. Describe the steps you would take to make 100 mL of a 0.200 mol/L sodium chloride solution from salt crystals. Include the equipment and calculations you would make. Remember this is not a reaction.
- 1) Calculate moles – $n=CV$; $n=0.200 \text{ mol/L} \times 0.100 \text{ L}$; $n=0.0200 \text{ mol}$ Na=22.99
 - 2) Calculate mass – $m=nM$; $m=0.0200 \text{ mol} \times 58.44 \text{ g/mol}$; $m=1.17 \text{ g}$ Cl=35.45
 - 3) Weight with a scale; Mix in beaker with 50 mL of water. 58.44 g/mol
 - 4) Place solution in a 100 mL volumetric flask and fill to the meniscus/calibration line
 - 5) Cap and mix

3. Describe the steps you would take to make 250 mL of a 0.453 mol/L solution of copper (II) sulphate from solid copper (II) sulphate pentahydrate. Include equipment and calculations.

1) Calculate the moles – $n=CV$; $n=0.453 \text{ mol/L} \times 0.250 \text{ L}$; $n=0.113 \dots \text{ mol}$ Cu=63.55

2) Calculate the mass – $m=nM$; $m=0.113 \dots \text{ mol} \times 249.71 \text{ g/mol}$; $m=28.3 \text{ g}$

3) Weight it; mix in beaker with about 125 mL

4) Place in a 250 mL volumetric flask and fill to the meniscus/calibration line

5) Cap and mix

Equipment: Calculator, weight scale, 250 mL volumetric flask, 125 mL beaker, eye dropper, cap

4. Describe the steps you would take to **make** 100 mL (V_2) of a 0.50 mol/L (C_2) sucrose solution from a container of 2.10 mol/L (C_1) sucrose solution. Include equipment and calculations.

1) Calculate volume that needs to be removed. $V_1=C_2V_2/C_1$;

$$V_1=0.50 \text{ mol/L} \times 0.100 \text{ L} / 2.10 \text{ mol/L}$$

$$V_1=0.0238 \text{ L}; V_1=24 \text{ mL}$$

2) Remove 24 mL with a graduated pipet

3) Place in a 100 mL volumetric flask; fill to calibration line; cap and mix

5. Describe the steps you would take to make 500 mL (V_2) of a 0.900 mol/L (C_2) sulphuric acid from a 1.50 L (V_1) container of 6.00 mol/L (C_1) sulphuric acid solution. Include equipment and calculations.

1) Calculate volume; $V_1=C_2V_2/C_1$; $V_1=0.900 \text{ mol/L} \times 0.500 \text{ L} / 6.00 \text{ mol/L} = 0.0750 \text{ L}$ or 75.0 mL

2) Remove 75 mL with a volumetric pipet.

3) Place in a 500 mL volumetric flask; fill with 425 mL of water to calibration line and cap and mix

6. What is the final concentration of a cleaner if 10 L(**V1**) of concentrated sodium hydroxide (19.1 mol/L)**C1**) is diluted to 400 L(**V2**)?
 $C_2 = C_1V_1/V_2$; $19.1\text{ mol/L} \times 10\text{ L} / 400\text{ L} = 0.48\text{ mol/L}$
7. What is the mass of baking soda (sodium hydrogen carbonate) needed to make 2.5 L of a 1.00 mol/L solution?
 1) **$n = CV$; $1.00\text{ mol/L} \times 2.5\text{ L}$; $n = 2.5\text{ mol}$**
 2) **$m = nM$; $2.5\text{ mol} \times 84.01\text{ g/mol}$; $m = 2.1 \times 10^2\text{ g}$ or 0.21 kg**
8. If 2.0 L of water is added to 1.0 L of a 0.250 mol/L solution of potassium hydroxide what is the final concentration. (Be Careful)
 $C_2 = C_1V_1/V_2$; $C_2 = 0.250\text{ mol/L} \times 1.0\text{ L} / 3.0\text{ L}$; $C_2 = 0.083\text{ mol/L}$
9. CHALLENGE: If 1.50 L of a 12.4 mol/L solution of hydrochloric acid was mixed with 300 mL of a 6.10 mol/L solution of hydrochloric acid, then what would be the final concentration.
 **$C_{\text{new}} = n_{\text{total}}/V_{\text{total}}$; $C_{\text{new}} = (1.50\text{ L} \times 12.4\text{ mol/L}) + (6.10\text{ mol/L} \times 0.300\text{ L}) / (1.50\text{ L} + 0.300\text{ L})$
 $C_{\text{new}} = (18.6\text{ mol} + 1.83\text{ mol}) / 1.8\text{ L}$; $C_{\text{new}} = 11.4\text{ mol/L}$**

10. CHALLENGE: How much water is added to 50.0 mL(**V1**) of a 0.500 mol/L(**C1**) solution to make a 0.100 mol/L(**C2**) solution?

$$\mathbf{V2 = C1V1/C2; V2 = 0.500\text{mol/L} \times 0.050\text{L} / 0.100\text{mol/L}; V2 = 250\text{ mL}}$$

$$\mathbf{V_{\text{water}} = V2 - V1; V_{\text{water}} = 250\text{ ml} - 50\text{ mL}; V_{\text{water}} = 200\text{ mL}}$$

Worksheet 2.4: Dissociation and ionization reactions

Name: _____

1. What type of compounds dissociate? What type of compounds ionize?

Ionic


Acids & molecular gases

2. Write dissociation or ionization reactions for the following chemicals after they are mixed with water. Show the physical states of all species involved. Use modified ionization reactions when necessary.


- a) Solid hydrochloric acid



- b) Solid strontium hydroxide




- c) Solid copper (II) sulphate pentahydrate



- d) Solid sodium bicarbonate (hydrogen carbonate)




- e) ammonia gas (acid and bases)



3. For each of the following write dissociation or ionization equations and find the concentration of each ion.


- a) 0.90 mol/L solution of sodium phosphate



$$0.90\text{mol/L}/1\text{mol} = x/3\text{mol} = x/1\text{mol}$$


$$x = 2.7\text{mol/L} \quad x = 0.90\text{mol/L}$$

- b) 0.143 mol/L solution of nitric acid



$$0.143\text{mol/L} \quad 0.143\text{mol/L}$$

- c) 0.0135 mol/L solution of calcium hydroxide



$$0.0135\text{mol/L}/1\text{mol} = x/1\text{mol} = x/2\text{mol}$$

$$x=0.0135 \text{ mol/L} \quad x=0.0270 \text{ mol/L}$$

d) 0.150 mol of hydrogen fluoride gas bubbled into 1.00 L of water
 $\text{HF}_{(g)} + \text{H}_2\text{O}_{(g)} \rightarrow \text{H}_3\text{O}^+_{(aq)} + \text{F}^-_{(aq)}$

0.150mol **0.150mol/L** **0.150mol/L**

4. What is the concentration of chloride ions in a solution prepared by dissolving 800 g of zinc chloride in 4.50 L of water?

1) $n=m/M$; $n=800/136.28\text{g/mol}$; $n=5.87\dots\text{mol}$



5.87 mol 11.74...mol

$C=n/V$; $C=2.61\text{mol/L}$

5. What is the mass of calcium chloride required to prepare 2.000 L of 0.120 mol/L chloride ions?

1) $n=CV$; $0.120\text{mol/L} \times 2.00\text{L} = 0.240\text{mol}$



3) $X/1\text{mol} = 0.240\text{mol}/2\text{mol}$

$X=0.120\text{mol}$

4) $m=nM$; $m=0.120\text{mol} \times 110.98\text{g/mol}$; 13.3 g




6. What is the final concentration if 2.0 L of water is added to 4.50 L of a 0.89 mol/L solution of sodium chloride?

$C_2=C_1V_1/V_2$; $C_2=0.89\text{mol/L} \times 4.50\text{L}/6.50\text{L}$; $C_2=0.62\text{mol/L}$



Worksheet 4.5: Net Ionic Equations

For the following reactions, write the nonionic equation, the total ionic equation and the net ionic equation.

1. Aqueous solutions of sodium sulphate and barium bromide are mixed.






2. A lead (II) nitrate solution reacts with sodium sulphide solution





NET IONIC:

3. Sulphuric acid is neutralized by a potassium hydroxide solution






4. Hydrochloric acid is added to a solution of barium hydroxide

5. Magnesium metal is added to an aqueous solution of hydrogen bromide





6. Zinc reacts with copper (II) sulphate solution

7. Zinc reacts with acetic acid (vinegar)



8. Bromine is added to a magnesium iodide solution



Worksheet 2.6: Solution Stoichiometry

1. A 200 mL solution of potassium phosphate reacts with 100 mL of 0.150 mol/L iron (III) sulphate solution. What is the concentration of the potassium phosphate solution?

2. If 230 mL of a 1.00 mol/L solution of aluminium chlorate is reacted with sufficient lithium hydroxide solution, what mass of precipitate is formed?



$$0.230\text{L} = V$$

$$1.00 \text{ mol/L} = C$$

$$n = CV; n = 1.00 \text{ mol/L} \times 0.230\text{L}$$

$$0.230 \text{ mol} / 1 \text{ mol}$$

$$x / 1 \text{ mol}$$


$$x = 0.230 \text{ mol}$$

$$m = nM; m = 0.230 \text{ mol} \times 78.01 \text{ g/mol}$$

$$m = 17.9 \text{ g}$$

3. Predict the mass of magnesium metal that will be required to react with 44.0 ml of 0.200 mol/L hydrochloric acid.

4. What volume of 1.00 mol/L $\text{HNO}_{3(aq)}$ is required to react completely with 1.20g of $\text{LiOH}_{(aq)}$?



2) $V=?$ $n=1.20\text{g}/23.95\text{g/mol}$

V $n=0.050104\text{mol}$

3) $X/1\text{mol}$ $= 0.050104\dots\text{mol}/1\text{mol}$


$X=0.050104\dots\text{mol}$

4) $V=n/C$; $V=0.050104\dots\text{mol}/1.00\text{mol/L}$

$V=0.0501\text{L}$ or 50.1 mL

5. A 100 ml sample of sodium sulphide solution is completely reacted with 50.0 ml of 0.250 mol/L lead (II) nitrate solution. Predict the concentration of the $\text{Na}_2\text{S}_{(\text{aq})}$?

6. 500 ml of 0.150 mol/L cobalt (II) nitrate solution is reacted with 500 ml of 0.250 mol/L of sodium hydroxide solution producing 4.77 g of precipitate. Find the % yield for this reaction.



$$0.075 \text{ mol} / 1 \text{ mol} =$$

$$x / 2 \text{ mol}$$

$$x = 0.150 \text{ mol}$$

0.125 < 0.150 LIMITING

$$0.125 \text{ mol} / 2 \text{ mol} =$$


$$x / 1 \text{ mol}$$

$$x = 0.0625 \text{ (5.809..g)}$$

$$\% \text{ yield} = A/T \times 100; \quad \% \text{ yield} = 0.0513 \dots \text{ mol} / 0.0625 \text{ mol} \times 100; \quad \% \text{ yield} = 82.1\%$$



7. CHALLENGE: Predict the final mass of a 500 g bar of lead that is allowed to react completely with 2.00 L of 2.00 mol/L HCl.



$$500 \text{ g}$$

$$n = CV$$

$$n = 2.00 \text{ mol/L} \times 2.00 \text{ L}$$

$$n = 4.00 \text{ mol}$$

$$x / 1 \text{ mol} =$$


$$4.00 \text{ mol} / 2 \text{ mol}$$

$$x = 2.00 \text{ mol}$$

$$m=nM; m=2.00\text{mol} \times 207.19\text{g/mol}; m=414.38\text{ g}$$

$$m_{\text{final}} = 500 - 414.38 = 85.6\text{ g}$$

8. A 75.0 mL sample of 0.25 mol/L silver chlorate solution reacts with 19.0 mL of 0.50 mol/L copper (II) sulphate solution. What is the concentration of the solution produced? (NOTE: Find out what the total volume of the solution produced.)



$$2) \quad n=CV \quad \quad \quad n=CV \quad \quad \quad C=?$$

$$n=0.25\text{mol/L} \times 0.075\text{L} \quad \quad \quad n=0.50\text{mol/L} \times 0.019\text{L}$$

$$n=0.01875\text{ mol} \quad \quad \quad n=0.0095\text{ mol}$$

$$0.01875/2\text{mol} \quad \quad \quad = \quad \quad \quad x/1\text{mol}$$

$$x=0.009375\text{...mol}$$

$$\underline{0.0095 > 0.009375\text{...mol EXCESS}}$$

$$0.01875/2\text{mol} \quad \quad \quad = \quad \quad \quad x/1\text{mol}$$

$$x=0.009375\text{...mol}$$

$$C=0.009375\text{mol}/0.094\text{L}$$

$$C=n/V;$$

$$C=0.0997; \quad C=0.10\text{ mol/L}$$

Worksheet 2.7: Review of Solutions

1. Answer the following questions

a) How do solutions differ from heterogeneous mixtures?

Solutions are uniform and appear as one substance – heterogeneous do not.

b) How do the number of molecules of $C_{12}H_{22}O_{11}$ in 250 mL of a 1.5 mol/L solution of $C_{12}H_{22}O_{11}$ compare to the number of molecules of $C_6H_{12}O_6$ in 250 mL of a 1.5 mol/L $C_6H_{12}O_6$?


The number of molecules is the same ($n=CV$); the mass is different

c) What is the term used to describe two liquids which will **NOT** mix with each other?
immiscible


d) What are two factors that affect the amount of solute that dissolves and two factors that affect the rate of dissolving? **Amount: temperature, pressure Rate: temperature, surface area, agitation**

2. Write the equation for each of the following dissolving in water. Use modified Arrhenius theory.


a) Hydrogen chloride gas




b) Solid aluminum nitrate



c) Solid sucrose



d) Aqueous nitric acid



3. Determine the concentration of each of the following solutes in the solution described.

a) 0.725 mol of cobalt (II) nitrate in 1.35 L of solution.

$$C=n/V; C=0.725\text{mol}/1.35\text{L}; C=0.537 \text{ mol/L}$$

b) 15.0 g of barium sulphate in 125 mL of solution.

$$n=m/M; n=15.0\text{g}/233.39\text{g/mol}; n=0.0642\dots\text{mol}$$

Ba=137.33

S=32.06

$$C=n/V; C=0.065\dots\text{mol}/0.125\text{L}; C=0.514\text{mol/L}$$


$$Ox4=64.00/233.39\text{g/mol}$$

c) 1.85×10^{22} molecules of ammonia gas in 400 mL of solution.

$$n=p/P; n=1.85 \times 10^{22} / 6.02 \times 10^{23}; n=0.0307\dots\text{mol}$$

$$C=n/V; C=0.0307\dots\text{mol}/0.400\text{L}; C=0.0768 \text{ mol/L}$$


4. Write the dissociation equation and calculate the concentration of each of the ions produced in 1.25 mol/L solution of barium chloride.



$$1.25\text{mol/L}/1\text{mol} = x/1\text{mol} = x/2\text{mol}$$

$$x=1.25 \text{ mol/L} \quad x=2.50 \text{ mol/L}$$



5. Write the dissociation equation and determine the concentration of the solution if 1.26 mol/L of $[Na^+]$ is found in a sodium phosphate solution.



$$x/1\text{mol} = 1.26\text{ mol/L}/3\text{mol}$$

$$x = 0.420\text{ mol/L}$$

6. CHALLENGE: What is the [Cl⁻] in a solution made by mixing 200 mL of 0.300 mol/L sodium chloride solution with 350 mL of 0.250 mol/L calcium chloride solution?



$$C = n_{\text{total}}/V_{\text{total}}$$

$$n = CV; n = 0.300\text{ mol/L} \times 0.200\text{ L}$$

$$n = 0.0600\text{ mol}/1\text{ mol} = x/1\text{ mol}$$

$$x = 0.0600\text{ mol}$$

$$n = CV; n = 0.250\text{ mol/L} \times 0.350\text{ L}$$

$$n = 0.0875\text{ mol}/1\text{ mol} = x/2\text{ mol}$$


$$x = 0.175\text{ mol}$$

$$C = 0.235\text{ mol}/0.550\text{ L}$$


$$C = 0.427\text{ mol/L}$$

Write net ionic equations for the following reactions. (3 marks)

- a) lead nitrate solution is mixed with sodium hydroxide




- b) barium nitrate reacts with potassium sulphide




NO NET IONIC EQUATION

- c) nitric acid reacts with barium hydroxide



7. Draw a diagram describing how methanol is dissolved in water. (1 mark)



8. Predict whether the following solutes are electrolytes or nonelectrolytes:

- nitrogen monoxide – **nonelectrolyte (molecular)**
- hydrofluoric acid – **electrolyte (acid)**
- magnesium hydroxide – **nonelectrolyte (ionic BUT not aqueous)**
- potassium hydrogen carbonate – **electrolyte (ionic & aqueous)**

9. A scientist wants to make 100 mL of a 0.150 mol/L sodium hydroxide solution. He has 100 g of solid sodium hydroxide and he has 1.00 L of a 2.25 mol/L sodium hydroxide solution. Describe step by step the two ways that he could make his 0.150 mol/L solution. Include the sample calculations and equipment.

Method I - from solid

1) Find moles; $n = CV$; $0.150 \times 0.100 = 0.0150$

$V_1 = C_2 V_2 / C_1 = 0.150 \times 0.100 / 2.25 = 6.67 \text{ mL}$

2) Find mass: $m = nM$; $0.0150 \times 40.00 = 0.600 \text{ g}$

3) Weigh on a scale
line

4) Mix 0.600g in 50 ml of water

5) Place in 100 mL volumetric flask
and fill to calibration line. Cap and mix

Method II - dilution

1) Find volume;

2) Remove 6.67mL with a graduated pipet

3) Place in a 100 mL volumetric flask and fill to

4) Cap and mix

10. A 20.0 g sample of lead (II) nitrate is mixed in 1.00 L of water. The lead (II) nitrate solution then reacts with a 1.00 L of a 0.100 mol/L solution of rubidium iodide. If 20.0 g of precipitate forms, what is the percent yield?

Worksheet 2.8: Introduction to Acids & Bases

1. Safety is very important when working with acids. Describe what the student should do in the following situations.
- A student drops a 100 mL beaker with 50 mL of hydrochloric acid and spills the acid onto the floor.
Report the accident to a teacher. Place baking soda until it stops bubbling. (If you have no baking soda, dilute the acid with water.)
 - A student drips a couple of drops of sodium hydroxide solution onto his hand.
Report the incident to a teacher. Wash the sodium hydroxide off with cold water.
 - A beaker with $\text{Ba}(\text{OH})_2$ tips over onto the lab bench.
Report the incident to a teacher. Place vinegar (acetic/ethanoic acid) onto the base. (If you have no vinegar, dilute the base with water.)
 - A student would like to dilute an acid and would like to know if he should add the acid to the water or the water to the acid
Always add acid to water (A to W).

2. WHMIS symbols help communicate dangers.

- WHMIS stands for **Workplace Hazardous Materials Information System**
- The symbol that would be associated with a beaker of base that corrodes metal is



Corrosive material

- Acids and bases can cause immediate and serious damage to a person's skin. The WHMIS symbol related to this is



Material causing immediate and serious toxic effect

- Some acids react with oxygen. The WHMIS symbol found on a bottle of this acid would be



Oxidizing Material

3. A person would like to make 100 mL 1.00 mol/L solution of NaOH. Describe the steps the student would use. Include the calculations.

- 1) Calculate the moles: $n = CV$; $n = 1.00 \text{ mol/L} \times 0.100 \text{ L}$; $n = 0.100 \text{ mol}$
- 2) Calculate mass: $m = nM$; $m = 0.100 \text{ mol} \times 40.00 \text{ g/mol}$; $m = 4.00 \text{ g}$
- 3) Weigh with scale; mix in beaker with 50 mL of water.
- 4) Place in 100 mL volumetric flask, fill to calibration line, cap & mix.

4. A person would like to dilute a 12.1 mol/L solution of HCl and make a 250 mL 3.00 mol/L solution. Describe the steps the student would use. Include the calculations.

- 1) Find the volume; $v_1 = C_2V_2/C_1$; $V_1 = 3.00 \text{ mol/L} \times 0.250 \text{ L} / 12.1 \text{ mol/L}$; $V = 62.4 \text{ mL}$
- 2) Remove it with graduated pipet.
- 3) Place in 250 mL volumetric flask; fill to calibration line; cap & mix

5. Indicators change color to indicate whether you have an acid or base. Litmus paper and bromothymol blue are two common indicators. Complete the following table for these indicators.

<u>PH</u>	<u>Litmus Paper color</u>	<u>Bromothymol Blue color</u>
3	Red	yellow
7	No change	Green
10	Blue	Blue

6. What is one property that is similar between acids and bases?


Both electrolytes, both dissolve in water (aqueous)

7. What is one property that is different between acids and bases?

PH, taste, touch, reactions


8. Complete the following acid or base reactions.

- a) sulphuric acid is neutralized by potassium hydroxide. Identify the "salt" in the reaction.




salt

b) hydrochloric acid reacts with magnesium



salt

c) self ionization of water



No salt

Worksheet 2.9: Acid & Base Calculations

1. A 1.00 L solution of 1.50 mol/L perchloric acid is diluted by adding 500 mL *of water*. What is the hydronium concentration of the diluted solution? $V_2 = V_1 + V_{\text{water}} = 1.00\text{L} + 0.500\text{L}$




$$C_2 = C_1 V_1 / V_2$$

$$C_2 = 1.50 \text{ mol/L} \times 1.00 \text{ L} / 1.50 \text{ L}$$

$$C_2 = 1.00 \text{ mol/L} \quad 1.00 \text{ mol/L} / 1 \text{ mol} = X / 1 \text{ mol}$$

$$X = 1.00 \text{ mol/L}$$

2. A 250 mL solution of 3.56 mol/L barium hydroxide is sitting on the counter in the lab. Help a chemistry 20 student determine the hydronium concentration of the solution.



$$n = CV; n = 3.56 \times 0.250 = 0.89 \text{ mol}$$


$$X \text{ 2 mol} / 1 \text{ mol} = 1.78 \text{ mol}$$

$$C = n/V; C = 1.78 \text{ mol} / 0.250 \text{ L} = 7.12 \text{ mol/L}$$

$$\text{H}_3\text{O}^+ = K_w / [\text{OH}^-]; \text{H}_3\text{O}^+ = 1 \times 10^{-14} / 7.12$$

$$= 1.40 \times 10^{-15} \text{ mol/L}$$

3. A 1.00 mol/L solution of nitric acid ionizes. What is the hydroxide ion concentration?



$$1.00 \text{ mol/L}$$

$$X / 1 \text{ mol}; X = 1.00 \text{ mol/L}$$

$$[\text{OH}^-] = K_w / [\text{H}_3\text{O}^+_{(aq)}]$$

$$= 1.00 \times 10^{-14} (\text{mol/L})^2 / 1.00 \text{ mol/L}; [\text{OH}^-] = 1.00 \times 10^{-14} \text{ mol/L}$$

4. A student takes 11.6 grams of strontium hydroxide and adds it to 3.00 litres of water. What is the hydronium concentration?

5. A solution contains 1.67×10^{-14} mol/L of hydronium ions. Determine the mass of barium hydroxide that was added to 1.00 L of water to make this solution.



$$[\text{OH}^-] = K_w / [\text{H}_3\text{O}^+_{(aq)}]$$

$$= 1.00 \times 10^{-14} (\text{mol/L})^2 / 1.67 \times 10^{-14} \text{ mol/L}$$

$$X \text{ 1 mol} / 2 \text{ mol}$$

$$0.5988 \dots \text{ mol/L}$$

$$X = 0.2994 \dots \text{ mol/L}$$


$$n = CV; n = 0.2994 \times 1 \text{ L} = 0.2994 \dots \text{ mol}$$

$$m = nM; m = 0.2994 \dots \text{ mol} \times 171.35 \text{ g/mol}$$

$$m = 51.3 \text{ g}$$


6. What is the concentration of hydroxide ions found in a 1.00 L solution of 2.00 mol/L potassium hydroxide?

7. What is the hydroxide concentration of a 1.00 L solution of 2.50 mol/L hydrobromic acid?



8. What is the hydronium concentration when 1.00 mol/L of barium hydroxide dissociates


9. 6.02×10^{22} particles of sulphuric acid ionize into hydrogen sulphate ions in 1.00 L of water. What is the hydroxide concentration of the solution?



10. A solution contains 3.45×10^{-12} mol/L of hydroxide ions. What is the concentration of the hydrochloric acid solution that contain these hydroxide ions?

Worksheet 2.10: Acid & Base Review

1. The concentration of hydroiodic acid is 1.73×10^{-3} mol/L. What is the pH and the pOH?



$$1.73 \times 10^{-3} \text{ mol/L/1mol} = X/1\text{mol}$$

$$X = 1.73 \times 10^{-3} \text{ mol/L}$$

$$\text{PH} = -\log (1.73 \times 10^{-3} \text{ mol/L})$$

$$\text{PH} = 2.76195\dots (2.762)$$

$$\text{POH} = 14 - \text{pH} = 11.238$$

2. What is the hydronium concentration and hydroxide concentration of a 2.47×10^{-2} mol/L solution of thallium hydroxide?

3. Complete the following table (Significant digits are important):

pH	[H ⁺] or [H ₃ O ⁺]	[OH ⁻]	pOH	A/B/N
14 -4.56 OR -log(3.6E-10) = 9.44	1E-14/2.8E-5 =3.6E-10mol/L OR 10^{-9.44}	=10^{-4.56} = 2.8E- 5mol/L OR 1E-14/3.6E-5	4.56	B
2) 14-4.910 =9.090	8.13 x 10⁻¹⁰ mol/L	1.23 X 10 ⁻⁵	-log (1.23E-5) =4.910	B
3) 7.449	3.56 x 10 ⁻⁸	2.81 x 10⁻⁷mol/L	6.551	B
4) 12.8	2 x 10⁻¹³mol/L	6. x 10⁻²mol/L	1.2	B
5) 3.52	3.0 x 10⁻⁴	3.3 x 10⁻¹¹	10.48	A
6) 13.759	1.74 x 10⁻¹⁴	5.74 X 10 ⁻¹	0.241	B

pH	[H ⁺] or [H ₃ O ⁺]	[OH ⁻]	pOH	A/B/N
7) 6.55	2.8×10^{-7}	3.5×10^{-8}	7.45	A
8) 2.399	3.99×10^{-3}	2.51×10^{-12}	11.601	A
9) 12.77	1.7×10^{-13}	5.9×10^{-2}	1.23	B
10) 5.95	1.1×10^{-6}	8.9×10^{-9}	8.05	A

4. What color would the indicator be given the following data:

	ORANGE IV	METHLY RED	PHENOL RED	METHYL ORANGE	INDIGO CARMINE
pOH=9.00 pH = 5.00	yellow	Red + Yellow = orange	yellow	yellow	blue
pH=8.3	Yellow	Yellow	Red	Yellow	Blue
[H+]=9.5 x 10 ⁻⁴ pH = 3.02	Yellow	Red	Yellow	Red	Blue
[OH-]=5.6 x 10 ⁻³ pOH = 2.25; pH = 11.75	Yellow	yellow	Red	Yellow	Blue + yellow = green
[H ₃ O+] = 1.0 x 10 ⁻⁷	Yellow	Yellow	Yellow to red = orange	Yellow	Blue

